

Investigating the Role of Different Factors in the Quality of Professional Life of Nurses: A Review Study

Shaher Hamaideh^{1*}, Abdallah Abu Khait¹, Hanan Al-Modallal¹, Rami Masa'deh²,
Ayman Hamdan-Mansour³, Mohammed AlBashtawy⁴

¹*Department of Community and Mental Health Nursing, Faculty of Nursing, Hashemite University, Zarqa 13133, Jordan.*

²*School of Nursing, Applied Science Private University, Amman 11937, Jordan.*

³*Department of Community Health, School of Nursing, The University of Jordan, Amman 11942, Jordan.*

⁴*Princess Salma Faculty of Nursing, Al al-Bayt University, Mafrq 25113, Jordan.*

Abstract

Nurses are one of the important pillars in health care systems and providing care to patients. Therefore, the quality of their professional life has a complex relationship with the characteristics of the work environment, personal life environment, and exposure to occupational accidents. Therefore, the present study was conducted to investigate the quality of professional life of nurses and the factors affecting it by reviewing past studies. For this purpose, using the keywords of health, public health, quality of professional life, and nurses, it was investigated in Pubmed, Proquest, Google Scholar, Scopus, and Web of Science databases. The results of various studies showed that the quality of professional life of nurses is average. In some studies, the quality of nurses' professional lives has been reported as optimal. In addition, the results of the studies showed that the quality of professional life in nurses is influenced by various factors such as salary fairness, justice, job satisfaction, desire to leave the profession, high workload, increased overtime, fatigue, inappropriate rules, decision-making power, Empathic communication, and occupational stress. Considering the average level of the quality of professional life of nurses, it is necessary to pay attention to the factors affecting the quality of professional life to maintain and improve the quality of nursing care.

Keywords: Quality of professional life, Nurses, Health, Public health

Introduction

Nursing is one of the main fields of care and a healthy nurse is one of the main factors to improve the quality of care. There is a positive and strong relationship between healthy nursing and health promotion because nurses are potential role models for society [1-3]. The World Health Organization defines health as not only the absence of disease but also the integration and harmony of well-being between physical, mental, social, and spiritual factors [4, 5]. Health, which has different physical, mental, and social dimensions, is one of the necessary conditions for fulfilling individual and social roles, and people can be fully active if they feel healthy and their society feels healthy. Undoubtedly, someone with a health disorder cannot take on personal, family, and social obligations and requirements and he will be unable to do it properly [6-8]. People's jobs also affect their health. Some jobs threaten health more than other jobs by exposing a person to numerous and different physical, physical, psychological, and social stressors. Nursing is one of these jobs. Nursing has been proposed as one of the most dangerous and stressful jobs and one of the first four stressful professions in the world [9-11].

Corresponding author: Shaher Hamaideh
Address: Department of Community and Mental Health Nursing,
Faculty of Nursing, Hashemite University, Zarqa 13133, Jordan.
E-mail: ✉ shaher29@hu.edu.jo
Received: 28 August 2024; **Revised:** 28 November 2024;
Accepted: 04 December 2024

How to Cite This Article: Hamaideh S, Abu Khait A, Al-Modallal H, Masa'deh R, Hamdan-Mansour A, AlBashtawy M. Investigating the Role of Different Factors in the Quality of Professional Life of Nurses: A Review Study. J Integr Nurs Palliat Care. 2025;6:12-7. <https://doi.org/10.51847/k8xYY6EKKp>

The World Health Organization, after studying stressful jobs, announced that out of 130 jobs surveyed, nurses ranked 27th in going to the doctor in terms of health-related problems [6, 12]. Long working hours, lack of support, heavy workload, violence from patients, insufficient personnel, insufficient resources, lack of respect, lack of involvement in decisions, and lack of adequate equipment are some of the things that affect the health of nurses [13, 14]. The results of studies show that nurses suffer from all kinds of diseases such as musculoskeletal, cardiovascular, respiratory, infectious, parasitic, contagious, mental, and behavioral disorders. These diseases are more common in female nurses than in males, which can be caused by the high proportion of female nurses compared to males. The results also show that moving, lifting, and transferring patients cause one-third of musculoskeletal diseases in nurses. Standing for a long time also causes other problems such as leg pain and varicose veins. High expectations from nurses cause psychological diseases such as stress, anxiety, and insufficient sleep in nurses, as a result, nurses' sensitivity to pain increases and they feel more pain [9]. Nurses who do not have good general health; will not be able to provide good care such as physical and psychological support to the patients, and this increases the risk of mistakes and occupational accidents, the consequences of which will affect the patient and the nurse [15, 16]. In addition, health problems in nurses increase absenteeism; as a result, it affects both the quality of care and is costly for the organization [17].

As mentioned, general health is very important in nurses; because their health is at risk due to their profession and special conditions. In addition, the health level of nurses has a direct impact on how they perform. Therefore, any disorder in this group and their performance will have a significant impact on care and treatment [18].

Nurses are one of the important pillars in health care systems and providing care to patients. Therefore, the quality of their professional life has a complex relationship with the characteristics of the work environment, personal life environment, and exposure to occupational accidents. Therefore, the present study was conducted to investigate the quality of professional life of nurses and the factors affecting it by reviewing past studies.

Materials and Methods

This research was a review study that was conducted in 2023, during which the general health of nurses was examined. For this purpose, using the keywords of health, public health, quality of professional life, and nurses, it was investigated in Pubmed, Proquest, Google Scholar, Scopus, and Web of Science databases. MESH was also used to determine keywords. Two experts in the field of nursing determined keywords and these two experts did searching for phrases in databases. Then, one of the research colleagues checked and re-searched the sources and databases to ensure the adequacy of the search for information and articles. The criteria for entering the research included that the articles were published in the period from 2011 to 2023, the subject of the study was the general health of nurses working in medical centers, and the full text was easily available in reputable research and academic journals. Exclusion criteria included studies that only addressed nurses' health from one perspective, for example, only physical or mental health, studies that were unrelated to the topic, as well as articles that were only conducted on nursing students.

To determine the compatibility of the articles with the research topic, first, the title and then the abstract of the articles were examined, and after the approval of the article, the topic was studied by one of the researchers for further examination of all the texts of the articles. The research subject, sample size and sampling method, statistical analysis, and study results were extracted from the selected articles by the responsible author, evaluated, and confirmed by other researchers, and disagreements were resolved through negotiation.

Results and Discussion

Lifestyle is one of the concerns of people in organizations. Nurses are not exempt from this as health defenders. The results show that less than 5% of nurses have a healthy lifestyle, which includes having a healthy diet, regular exercise, maintaining body weight, and not smoking [19]. Because exercise is one of the dimensions of lifestyle, the results of the studies show that nurses scored low in this dimension. In this regard, the results of a study conducted on American nurses showed that 72% of nurses did not participate in physical activities [19]. Among the factors affecting the lack of ideal physical activity in most nurses, chronic fatigue and reduced mobility resulting from shift work can be mentioned. Of course, the role of urbanization and car life cannot be ignored. Due to inactivity, we should also worry about overweight and obesity, which are prone to cardiovascular diseases and diabetes. Therefore, participating in physical activities is effective in preventing and reducing the risk of osteoporosis and can help manage risk factors such as high cholesterol, high blood pressure, and body weight [20, 21].

A healthy diet is another important aspect of lifestyle, as there are many challenges in eating habits. The results of Wang *et al.* study showed that shift work is related to abnormal eating behaviors. So sweet foods, snacks, and foods with low nutritional value are considered as options that are more available compared to healthy foods during working nights, and with the increase of more than four working nights per month, nutritional habits falsehood increases in nurses [22]. Of course, limited facilities for storing heating, and eating homemade food,

time barriers, distance to buy food, and the variable quality of food were also other problems in this field [23]. The results of studies show that 61% of American nurses have an unhealthy diet [19].

In terms of smoking, in this study, nurses are at a favorable level, while smoking patterns among nurses are different between countries. The results of a study in New Zealand showed that 13% of female nurses (32,682 people) were smokers, which was the highest prevalence of smoking among nurses working in the psychiatric department [24]. In Turkey, 45% of nurses were smokers [25]. Due to the hard work conditions of working nurses the stress during work hours and the role of fatigue, nurses may use ineffective coping mechanisms such as excessive alcohol consumption and smoking [26].

Stress management is another dimension of lifestyle. The findings of studies show that nurses suffer a lot of stress due to the nature of their job, which is one of the major inevitable consequences of job stress, job burnout [27, 28]. The results of the Kurnat-Thoma *et al.* [29] study showed that burnout affects the lifestyle of nurses. On the other hand, stress management leads to an increase in the quality of life, which is confirmed by the results of Tal *et al.* research [30]. Therefore, it is necessary to use comprehensive approaches such as cognitive training, mindfulness, yoga, meditation, prayer, massage, and touch therapy that help manage stress and health [31]. Lifestyle has many effects on nurses' organizational citizenship behavior so the incompatibility between people's lifestyles and their interests and activities will lead to the occurrence of anti-organizational citizenship behaviors. When nurses conclude that the hospital fulfills its obligations, the incidence of organizational citizenship behaviors will increase [32]. The results of Nesai *et al.* study showed that e-learning was significant for strengthening health-promoting behaviors in nurses and the lifestyle score increased [33]. On the other hand, choosing a healthy lifestyle for nurses has positive effects on the care of the patient population [34]. Nurses, who practice a healthy lifestyle, including daily physical activity, healthy diet, and proper sleep, can better prevent workplace injuries and mistakes related to fatigue [35]. It also increases their ability to deal with job stress [36]. Based on the obtained results, most of the studies have reported the general health status of nurses as inappropriate or at risk. The most common health disorders of nurses were mental disorders high levels of depression and functional disorders. In addition, the results of the studies show that there is a relationship between factors such as job stress, job satisfaction, burnout, family life, emotional and spiritual intelligence, and religious beliefs with the general health of nurses. The investigated intervention studies show that compliance with things such as group therapy, stress management, communication skills training, and emotional intelligence training has improved public health and reduced anxiety and stress in nurses [37-39]. The results of a study also showed that the use of zinc food supplements improves the physical health of nurses [40].

The results of the current research show that most of the studies reviewed about the public health of nurses have described the current situation and related factors and have provided solutions to improve the public health of nurses. Nurses are in direct contact with society and touch people's problems closely. Therefore, it has a serious responsibility, and this causes an increase in job burnout among nurses and, as a result, a drop in their physical and mental health [41]. However, in the conducted studies, different results have been obtained regarding the health level of nurses. For example, in a study, Niazi *et al.* investigated the relationship between general health and emotional intelligence of nurses and reported the level of health of nurses as favorable [42]. Similarly, Farsi *et al.* reported the general health level of 68% of nurses in the normal range [43]. This is even though Dehghankar *et al.* have mentioned the general health status of nurses as inadequate in their study [18]. However, in the study of Barzideh *et al.* under the title of dimensions of occupational stress and its relationship with the general health status of nurses, the general health status of most nurses has been reported as doubtful [44]. In a similar study, Zanganeh *et al.* investigated the level of job burnout and its relationship with the general health of nurses, and the level of the general health of nurses was mentioned as average [41]. Finally, the results of the present study show that the general health status of nurses is inappropriately mentioned in more than 62% of articles.

In an extensive study conducted by Fernandes *et al.* [45] between 2010 and 2011 in Brazil in 18 hospitals with 3,229 nurses, the researchers examined the health and working hours of nurses by gender. In this study, the level of health is categorized into three groups: good, normal, and poor. The results of this research show that 6.85% of female nurses have good health, 1.27% have normal health, and only 1.7% have poor health. In addition, 65.5% of male nurses had good health, 6.6% had average health, and only 8.6% had poor health [42].

The results of studies show that the most focus on general health disorders in nurses are mental-psychological problems, depression, and anxiety rather than physical. For example, in a study, Haseli *et al.* investigated the general health status and related factors in nurses working in the teaching hospitals of Shiraz University of Medical Sciences. The findings of his study showed that 59.5% of nurses suspected mental disorders and 12.7% suspected physical disorders [46]. In their study, Dehghankar *et al.* showed that 1.6% of nurses have physical problems, 1.4% have anxiety and insomnia problems, 1.4% have functional disorders, and 1.6% have depression [18]. The results of the above and other studies show that nurses mentally, and psychologically, they are exposed to serious harm in their work environment. The evaluation of the studies also shows that several factors are related to the general health of nurses. For example, Haseli *et al.* [46] showed in their study that there is a significant relationship between mental health, job satisfaction, and satisfaction with the field of study. In addition, in their study, Mashak *et al.* investigated the relationship between nurses' occupational stress and general health status. The results of his

study show that stress can have adverse effects on nurses' performance and their general health by causing burnout. Therefore, there is a relationship between stress and the general health of nurses [47].

In a study conducted by Sauer and McCoy [48] in the United States of America on 345 nurses to investigate the effect of harassment on the health of nurses, the researchers concluded that nurses who are harassed in the workplace have less physical and mental health [48]. This study and similar studies show the importance of the work environment and its effect on nurses' health. Therefore, it can be said that the interaction of several work factors affects the health of nurses, although the relationship between the variables is debatable.

In connection with the promotion of public health, interventions have also been carried out in nurses and useful results have been presented. In a study conducted by McElligott *et al.* [49] in the USA on 149 nurses, the researchers examined the effect of behavioral methods on health promotion in nurses. The results of this study show that strategies such as holding training classes, performing activities such as massage reflexology, and simulation, and providing programs for stress management increase health in nurses.

The results also show that nurses suffer more psychologically, physically, and socially. The conducted studies also provide measures to improve the health of nurses. Actions such as group therapy, stress management, communication skills training, use of some nutritional supplements, and emotional intelligence training have made a significant contribution to increasing the health of nurses. This requires managers and officials to pay more attention to this issue and its importance in the health system. Therefore, it is recommended that the authorities pay more attention to the health of employees, especially nurses, and be diligent in improving it so that they can improve the quality of their services accordingly.

Conclusion

The results of various studies showed that the quality of professional life of nurses is average. In some studies, the quality of nurses' professional lives has been reported as optimal. In addition, the results of the studies showed that the quality of professional life in nurses is influenced by various factors such as salary fairness, justice, job satisfaction, desire to leave the profession, high workload, increased overtime, fatigue, inappropriate rules, decision-making power, Empathic communication, and occupational stress. Considering the average level of the quality of professional life of nurses, it is necessary to pay attention to the factors affecting the quality of professional life to maintain and improve the quality of nursing care.

Since the results of the study in some dimensions of the nurses' lifestyle, such as exercise, diet, and stress management, were reported at a low level, therefore, designing a comprehensive program to promote physical activity, integrating exercise into their daily life, teaching healthy diet daily consumption of fruits and vegetables and training to deal with stress seem necessary. Therefore, the researcher hopes that the results of this research can be useful in understanding the problems of nurses regarding the improvement of their lifestyle, and after that, they can take steps to solve their problems with proper planning.

Acknowledgments: None

Conflict of interest: None

Financial support: None

Ethics statement: None

References

1. Gandoy-Crego M, Clemente M, Mayán-Santos JM, Espinosa P. Personal determinants of burnout in nursing staff at geriatric centers. *Arch Gerontol Geriatr.* 2009;48(2):246-9. doi:10.1016/j.archger.2008.01.016
2. Yousaf M, Khan MM, Paracha AT. Leading professionally diverse workgroups of healthcare professionals for improving quality of care. *J Organ Behav Res.* 2021;6(1-2021):106-19.
3. Alghamdi M, Alshehri M, Sindi M. Quality of life assessment and co-morbidities of maxillofacial surgery department patients undergoing iliac crest harvesting procedures. *Annals Dent Spec.* 2022;10(4):91-100.
4. Sadock BJ, Sadock VA, Ruiz P. Synopsis of psychiatry: behavioral sciences. *Clin Psychiatry.* 2003;9:632.
5. Malik M, Mahjabeen M, Rana S, Hussain A, Hashmi A. Quality of life and depression among postmenopausal women in Pakistan. *Arch Pharm Pract.* 2021;12(3):29-33.
6. Maghsoodi S, Hesabi M, Monfared A. General health and related factors in employed nurses in medical-educational centers in Rasht. *J Holist Nurs Midwifery.* 2015;25(1):63-72.
7. Salhi B. Effect of the mandatory adoption of IFRS on the quality of accounting information: case of France. *J Organ Behav Res.* 2021;6(2):152-60.

8. Wąsacz K, Chomyszyn-Gajewska M. Oral health related quality of life (OHRQoL) and associated factors in adult patients. *Ann Dent Spec.* 2022;10(1):7-12.
9. Raeissi P, Raeissi N, Shokouhandeh L. The relationship between nurses' mental health and working motivation. *Adv Nurs Midwifery.* 2015;24(86):35-42.
10. AlHamoudi SS, AlRashidi SN, AlHarbi HA, AlNaji NJ, AlMohammed MA, Ansari SH. Confidence in performing endodontic treatment and perception of the quality of endodontic education. *Arch Pharm Pract.* 2021;12(2):94-8.
11. Saltykova YA. Organizational behavior of steel market players in the context of a pandemic. *J Organ Behav Res.* 2021;6(1):71-9.
12. AlHussain BS, AlShalawi SS, AlQahtani MA, AlMeshari SZ, AlShehri HM, AlAmeer BM, et al. The quality of life and satisfaction after orthognathic surgery in Saudi Arabia. *Ann Dent Spec.* 2022;10(4):117-25.
13. Bambra C, Gibson M, Sowden AJ, Wright K, Whitehead M, Petticrew M. Working for health? Evidence from systematic reviews on the effects on health and health inequalities of organisational changes to the psychosocial work environment. *Prev Med.* 2009;48(5):454-61. doi:10.1016/j.ypmed.2008.12.018
14. Alghamdi RS, Alharbi TS, Alsubaie WR. Quality standards of histopathology laboratory and work facilities in a developed country. *Arch Pharm Pract.* 2021;12(1):90-7.
15. Soleimany M, Nasiri Ziba F, Kermani A, Hosseini F. Comparison of the general health and sleep quality of nurses in two groups of without shift and with shift working. *Iran J Nurs.* 2008;20(50):29-38.
16. Aelita K, Natalia Z, Roksolana K, Olena K. Student youth health culture in system managing the quality of higher education. *Arch Pharm Pract.* 2021;12(2):86-93.
17. Arab M, Movahed Kor E, Hosseini M. Effective factor analysis on health related absence among nurses of clinical wards in general hospitals affiliated to Tehran university of medical sciences. *J Hosp.* 2016;15(1):21-30.
18. Dehghankar L, Askarian Omran S, Hassandoost F, Abbas Pour S, Jourabchi Z, Rafiei H. General health of Iranian registered nurses: a cross sectional study. *Int J Novel Res Healthc.* 2016;3(2):105-8.
19. Priano SM, Hong OS, Chen JL. Lifestyles and health-related outcomes of U.S. hospital nurses: a systematic review. *Nurs Outlook.* 2018;66(1):66-76.
20. Xu X, Byles JE, Shi Z, Hall JJ. Evaluation of older Chinese people's macronutrient intake status: results from the China health and nutrition survey. *Br J Nutr.* 2015;113(1):159-71.
21. McCullough ML, Patel AV, Kushi LH, Patel R, Willett WC, Doyle C, et al. Following cancer prevention guidelines reduces risk of cancer, cardiovascular disease, and all-cause mortality. *Cancer Epidemiol Biomark Prev.* 2011;20(6):1089-97.
22. Wong H, Wong MC, Wong SY, Lee A. The association between shift duty and abnormal eating behavior among nurses working in a major hospital: a cross-sectional study. *Int J Nurs Stud.* 2010;47(8):1021-7.
23. Perry L, Hoban K, Ryan E. Food for thought: nurses are what they eat. In *Proceedings of the 4th Annual Worldwide Nursing Conference, Singapore 2016 Jul* (pp. 18-19).
24. Edwards R, Bowler T, Atkinson J, Wilson N. Low and declining cigarette smoking rates among doctors and nurses: 2006 New Zealand census data. *N Z Med J.* 2008;121(1284):43-51.
25. Sezer H, Guler N, Sezer RE. Smoking among nurses in Turkey: comparison with other countries. *J Health Popul Nutr.* 2007;25(1):107-11.
26. Hensel D. Relationships among nurses' professional self-concept, health, and lifestyles. *West J Nurs Res.* 2011;33(1):45-62.
27. Nguyen HTT, Kitaoka K, Sukigara M, Thai AL. Burnout study of clinical nurses in Vietnam: development of job burnout model based on Leiter and Maslach's theory. *Asian Nurs Res.* 2018;12(1):42-9.
28. Walker LO, Avant KC. *Strategies for theory construction in nursing.* Upper Saddle River, NJ: Pearson/Prentice Hall; 2005. p. 63-76.
29. Kurnat-Thoma E, El-Banna M, Oakcrum M, Tyroler J. Nurses' health promoting lifestyle behaviors in a community hospital. *Appl Nurs Res.* 2017;35:77-81.
30. Tel A, Tavassoli E, Rad GS, Zadeh DS. Investigating the health promoting lifestyle and its relationship with quality of life in graduate students of school of public health, Isfahan university of medical sciences. *J Health Syst Res.* 2011;7(4).
31. Markwell P, Polivka BJ, Morris K, Ryan C, Taylor A. Snack and relax®: a strategy to address nurses' professional quality of life. *J Holist Nurs.* 2016;34(1):80-90.
32. Ertürk A. Increasing organizational citizenship behaviors of Turkish academicians: mediating role of trust in supervisor on the relationship between organizational justice and citizenship behaviors. *J Manag Psychol.* 2007;22(3):257-70.
33. Tsai YC, Liu CH. An eHealth education intervention to promote healthy lifestyles among nurses. *Nurs Outlook.* 2015;63(3):245-54.
34. Zhu DQ, Norman IJ, While AE. The relationship between doctors' and nurses' own weight status and their weight management practices: a systematic review. *Obes Rev.* 2011;12(6):459-69.

35. Geiger-Brown J, Rogers VE, Trinkoff AM, Kane RL, Bausell RB, Scharf SM. Sleep, sleepiness, fatigue, and performance of 12-hour-shift nurses. *Chronobiol Int.* 2012;29(2):211-9.
36. Wright K. Alleviating stress in the workplace: advice for nurses. *Nurs Stand.* 2014;28(20):37-42.
37. Pashib M, Seyedmoharrami I, Mohammadi S, Maryam T. Stress management group therapy on job satisfaction and general health in nurses. *J Torbat Heydariyeh Univ Med Sci.* 2015;3(2):1-7.
38. Amini M, Nouri A, Samavatyan H. Effect of communication skills training on general health of nurses. *Health Inf Manag.* 2013;10(1(29)):1-9.
39. Sharif F, Rezaie S, Keshavarzi S, Mansoori P, Ghadakpoor S. Teaching emotional intelligence to intensive care unit nurses and their general health: a randomized clinical trial. *Int J Occup Environ Med.* 2013;4(3):141-8.
40. Gholipour Baradari A, Mahdavi A, Shokri Afra G, Hoseini SH, Yazdani Cherati J, Emadi SA, et al. Effect of zinc supplement on general health of ICU nurses. *J Mazandaran Univ Med Sci.* 2013;22(98):76-85.
41. Zanganeh S, Moradbeygi K, Rasteh M, Raisifar Z. Investigating the degree of job burnout and its relation with the general health of nurses working in hospitals of Abadan and Khoramshahr in 2012. *J Ilam Univ Med Sci.* 2014;22(Supplement):217-24.
42. Niazi M, Menati R, Delpisheh A, Menati S, Kassani A. The association between general health and emotional intelligence in the nurses of Ilam province. *Sadra Med J.* 2015;3(3):179-90.
43. Farsi Z, Dehghan Nayeri N, Sajadi A. General health status and burnout of the hematopoietic stem cell transplantation nurses in Tehran. *Int J Community Based Nurs Midwifery.* 2013;1(1):52-61.
44. Barzideh M, Choobineh A, Tabatabaei S. Job stress dimensions and their relationship to general health status in nurses. *Occup Med Q J.* 2013;4(3):17-27.
45. Fernandes JDC, Portela LF, Griep RH, Rotenberg L. Working hours and health in nurses of public hospitals according to gender. *Rev Saude Publica.* 2017;51(63):1-13. doi:10.1590/S1518-8787.2017051006808
46. Haseli N, Ghahramani L, Nazari M. General health status and its related factors in the nurses working in the educational hospitals of Shiraz university of medical sciences, Shiraz, Iran, 2011. *Nurs Midwifery Stud.* 2013;1(3):146-51.
47. Mashak B, Farhand B, Moghadam S, Pazhoom Z, Hajalikhani T, Taghipoor N, et al. Relationship between job stress among nurses with their general health status in Kamali hospital in 2013. *Alborz Univ Med J.* 2015;4(4):231-6.
48. Sauer PA, McCoy TP. Nurse bullying: impact on nurses' health. *West J Nurs Res.* 2017;39(12):1533-46. doi:10.1177/0193945916681278
49. McElligott D, Siemers S, Thomas L, Kohn N. Health promotion in nurses: is there a healthy nurse in the house? *Appl Nurs Res.* 2009;22(3):211-5.