

Management of the Airway by Paramedic Nurses in Out-of-Hospital Cardiac Arrest Situations

Akari Sato^{1*}, Naomi Fujii¹, Rina Takeda¹

¹Department of Nursing Leadership, Graduate School of Health, Osaka Metropolitan University, Osaka, Japan.

Abstract

Effective airway management is essential during out-of-hospital cardiac arrest (OHCA) to ensure adequate ventilation and oxygenation. Advanced airway techniques, including the use of a supraglottic airway device (SAD) or endotracheal tube (ETT), typically follow initial bag-valve-mask (BVM) ventilation, with emergency front-of-neck access reserved as a last-resort option. Over time, and particularly during the COVID-19 pandemic, EMS airway management protocols have evolved. This study aimed to evaluate how these protocol changes affected airway management practices and the success rates of different techniques. We performed an observational analysis using data from the ARREST registry spanning 2019–2023. All adult OHCA patients receiving resuscitation from EMS were included, while cases involving helicopter emergency medical services were excluded. Trends in the use of airway devices during resuscitation and their first-pass success rates were analyzed. The proportion of cases in which an SAD was used increased, while ETT usage declined. Specifically, SADs were used in 59% of cases (95% CI: 57–60%), ETTs in 21% (95% CI: 19–22%), and BVM alone in 14% (95% CI: 13–15%). First-pass success for ETT placement improved from 53% to 68%, whereas SAD first-pass success remained consistently high at 93%. During CPR performed by ambulance nurses, the use of SADs rose and ETT use declined. Although ETT first-pass success improved, it remained below guideline-recommended standards for prehospital intubation. SAD placement demonstrated high first-pass success, supporting current Dutch EMS guidelines that prioritize SADs as the primary advanced airway device during resuscitation. Nonetheless, further training and skill development in intubation are necessary for selected OHCA patients.

Keywords: Out-of-Hospital Cardiac Arrest (OHCA), Airway management, Supraglottic Airway Device (SAD), Endotracheal Intubation (ETT)

Introduction

Cardiopulmonary resuscitation (CPR) primarily aims to maintain blood oxygenation and circulation to vital organs, thereby preventing ischemic injury [1]. Establishing a patent airway is the first step in ensuring adequate oxygenation. Upon arrival at an OHCA scene, EMS personnel initiate chest compressions and BVM ventilation. While BVM ventilation can be rapidly applied, it requires continuous attention, and airway management can remain challenging even for experienced providers [2, 3]. Following BVM ventilation, advanced airway interventions can be attempted. Endotracheal intubation has historically been considered the “gold standard” during CPR, as it secures the airway and reduces gastric insufflation and aspiration [2, 4]. However, aspiration often occurs before EMS arrival and may still occur during or after intubation, limiting its effectiveness in aspiration prevention [2, 4].

Corresponding author: Akari Sato
Address: Department of Nursing Leadership, Graduate School of Health, Osaka Metropolitan University, Osaka, Japan.
E-mail: ✉ akari.sato.study@gmail.com
Received: 29 May 2021; **Revised:** 12 September 2021;
Accepted: 14 September 2021

How to Cite This Article: Sato A, Fujii N, Takeda R. Management of the Airway by Paramedic Nurses in Out-of-Hospital Cardiac Arrest Situations. *J Integr Nurs Palliat Care.* 2021;2:122-33. <https://doi.org/10.51847/DSZOrj87sO>

Successful first-pass intubation is linked to a higher likelihood of return of spontaneous circulation (ROSC) [5]. Conversely, providers with limited intubation experience may prolong interruptions to chest compressions, resulting in extended apnea periods and poorer survival outcomes [6, 7].

SADs offer an effective alternative for ventilating many patients and require less technical skill to insert and maintain proficiency [8–11]. Comparative studies have found that SADs may yield similar, or occasionally superior, survival outcomes relative to ETTs, prompting many national and international EMS protocols to shift toward SADs for OHCA advanced airway management [4, 12–17]. These shifts may influence provider experience and success rates with airway devices [18–25]. In the United States, there has been a trend toward decreased ETT use and increased SAD use, but the impact on first-pass success remains uncertain [26, 27].

This study aims to examine trends in airway management by Dutch EMS during CPR over time. Understanding the effects of protocol changes on airway strategy selection and performance can help identify potential needs for targeted training, alternative approaches, or device innovations.

Materials and Methods

Study setting

In the Netherlands, suspected out-of-hospital cardiac arrest (OHCA) cases prompt dispatch of two advanced life support (ALS) ambulances, each staffed with an ambulance nurse and a driver. Ambulance nurses are qualified to manage the airway using techniques such as bag-valve-mask (BVM) ventilation, supraglottic airway device (SAD) insertion, endotracheal intubation (ETT), and emergency front-of-neck access (eFONA). In parallel, volunteer responders and other first responders are mobilized to deliver early basic life support (BLS) and defibrillation [28, 29].

Certain OHCA scenarios trigger Helicopter Emergency Medical Services (HEMS) deployment [30]. This includes patients with suspected airway compromise, children (<18 years), pregnant patients, or those with arrest due to trauma, drowning, or electrocution. HEMS is also activated when ground EMS is expected to arrive after more than ten minutes. From October 2022, HEMS were additionally used in the ARREST study region for witnessed OHCA patients aged 18–50 as part of the ON-SCENE trial [31].

All ambulance nurses operate under national protocols consistent with the Dutch and European Resuscitation Council (ERC) guidelines [32]. The 2014 national protocol introduced SAD use and allowed providers to choose the advanced airway device based on clinical judgment or regional policy [15]. During the study period, no adjunct devices (e.g., video laryngoscopes or gum elastic bougies) were employed. COVID-19 temporarily modified these guidelines: between April and June 2020, EMS were instructed to skip BVM ventilation and place an SAD (primary) or ETT (secondary) [33]. From June 2020, these modifications applied only to suspected COVID-19 patients, and all pandemic-related changes were lifted by September 2021 [34, 35]. The direct SAD approach, bypassing BVM, remains uncommon outside of COVID-19-specific guidance. The updated 2023 guidelines, effective May 1, 2023, recommend ETT only when SAD is ineffective or unachievable [17], reflecting the 2021 ERC guidance that ETT should be attempted only in systems achieving $\geq 95\%$ success within two attempts [32].

Study design and data source

This observational study used the ARREST registry, a prospective database capturing all suspected OHCA events in North-Holland since 2005, covering roughly 2.7 million residents [36, 37]. ARREST systematically collects Utstein-recommended OHCA variables from multiple sources, including dispatch centers, volunteer responder systems, AED recordings, EMS defibrillators, EMS run sheets, EMS questionnaires, and hospital records [36, 38]. Survivors provide informed consent for full dataset inclusion; if consent is declined or unobtainable, a limited anonymized dataset is used. For deceased patients, full data inclusion is allowed without consent.

Airway-specific data collection started in 2019. HEMS dispatch and arrival were systematically recorded from October 2022 onward; for earlier cases, ARREST data were retrospectively matched with HEMS records to confirm HEMS involvement during CPR. Approximately 15% of pre-October 2022 patients (~1034) had anonymized data preventing matching; these cases were retained because the expected number affected by HEMS (~66) was small.

Ambulance nurses reported airway management details via EMS questionnaires, including the order of successfully placed devices, first-pass success for ETT and SAD, and the final airway device used. First-pass success was defined as achieving effective ventilation on the initial attempt. Subsequent successful placements following an unsuccessful first attempt were recorded in sequence. The definitive airway was the device in place at prehospital CPR termination or at emergency department arrival during ongoing CPR.

The study did not fall under the Medical Research Involving Human Subjects Act, as confirmed by the Amsterdam UMC Medical Ethics Review Board (W17_089).

Patient selection

All consecutive adult patients (aged ≥ 18 years) who experienced OHCA and received attempted resuscitation by EMS were considered for inclusion. Resuscitation by EMS was defined as either performing at least one two-minute cycle of CPR or delivering at least one shock using an EMS defibrillator. Patients were excluded if HEMS was primarily dispatched due to the cause of cardiac arrest or if HEMS arrived secondarily and were documented on scene, as they might have performed airway interventions.

Objectives

The primary objective was to determine, by year from 2019 to 2023, the proportion of adult OHCA patients receiving BVM, SAD, ETT, or eFONA as the definitive airway during CPR. A secondary objective was to compare the first-pass success rates of SAD and ETT placement during resuscitation, also by year. This study was not intended to evaluate the impact of airway strategies on patient outcomes.

Sample size considerations

No prospective sample size calculation was performed, as the cohort size was determined by the available cases in the registry during the study period. Post hoc, we assessed the study's ability to detect a clinically meaningful 5% change in the use of specific airway devices over five years [39]. With 5,222 patients analyzed, the study had 80% power to identify a 5.6% increase or decrease (two-sided) at an alpha of 0.05.

Statistical analysis

Sample size feasibility was assessed using Stata/BE 17.0 (StataCorp LLC). Statistical analyses were conducted with SPSS® version 28.0 (IBM® SPSS®, Chicago, IL). Categorical variables were summarized as percentages, while continuous variables were expressed as means with standard deviations or medians with interquartile ranges. Temporal trends in airway device use were evaluated using chi-square tests for trend. Proportions for the primary outcome were reported with 95% confidence intervals (CI). Differences between airway groups for categorical data were tested using chi-square tests; when expected cell counts were ≤ 5 , Fisher's exact test (for two groups) or Fisher-Freeman-Halton exact test (for multiple groups) was applied. Continuous variables across airway groups were compared using Kruskal-Wallis tests.

Due to the very low number of patients with eFONA ($n=4$), these cases were excluded post hoc to reduce the risk of patient re-identification.

Results and Discussion

Between 2019 and 2023, 13,098 suspected OHCA cases were recorded. Resuscitation was attempted by EMS in 6,248 cases. After excluding pediatric cases, instances where HEMS was primarily or secondarily present, and eFONA cases, 5,222 patients were included for analysis (**Figure 1**). Among these, 59 percent (95 percent CI: 57–60 percent) received an SAD as the definitive airway, 14 percent (95 percent CI: 13–15 percent) had BVM ventilation, and 21 percent (95 percent CI: 19–22 percent) underwent ETT placement by EMS at the conclusion of CPR (**Table 1**).




Figure 1. Patient inclusion flowchart. CPR= cardiopulmonary resuscitation; eFONA= emergency front-of-neck access; EMS= Emergency Medical Services; HEMS= Helicopter Emergency Medical Services; OHCA= out-of-hospital cardiac arrest. *Cases where return of spontaneous circulation (ROSC) occurred due to successful defibrillation by an AED prior to EMS arrival

Table 1. Patient baseline characteristics according to the type of definitive airway used.

	ETT	SAD	BVM	Unknown	Statistics	Total
Adult OHCA with CPR attempt, <i>n</i> (%; 95% CI)	1072 (21; 19–22)	3062 (59; 57–60)	711 (14; 13–15)	377 (7; 7–8)		5222
Male sex, <i>n</i> (%)	742 (69)	2130 (70)	521 (73)	267 (71)	<i>p</i> = 0.225	3660 (70)
Missing	0	1 (0)	0	1 (0)		2 (0)
Age in years, Median (IQR)	72 (62–80)	70 (60–78)	67 (55 – 76)	71 (60–79)	<i>p</i> < 0.001	70 (60–78)
First rhythm, <i>n</i> (%)					<i>p</i> < 0.001	
Shockable (VF/VT)	340 (32)	873 (29)	301 (42)	102 (27)		1616 (31)
PEA	329 (31)	978 (32)	116 (16)	90 (24)		1513 (29)
Asystole	291 (27)	910 (30)	119 (17)	104 (28)		1424 (27)
EMS Witnessed*	71 (7)	226 (7)	149 (21)	55 (15)		501 (10)
Unknown**	41 (4)	75 (2)	26 (4)	26 (7)		168 (3)
Cause of arrest, <i>n</i> (%)					<i>p</i> = 0.347	
Medical	1063 (99)	3032 (99)	705 (99)	370 (98)		5170 (99)
Drug overdose	9 (1)	30 (1)	6 (1)	7 (2)		52 (1)
Arrest witness, <i>n</i> (%)					<i>p</i> < 0.001	
Bystander	676 (63)	1773 (58)	372 (52)	161 (43)		2982 (57)
EMS	71 (7)	229 (8)	151 (21)	57 (15)		508 (10)
None	312 (29)	1033 (34)	178 (25)	128 (34)		1651 (32)
Missing	13 (1)	27 (1)	10 (1)	31 (8)		81 (2)
BLS before EMS arrival, <i>n</i> (%)	865 (81)	2470 (81)	564 (79)	291 (77)	<i>p</i> = 0.812	4190 (80)
Missing	5 (1)	16 (1)	3 (0)	21 (6)		45 (1)
AED before ambulance arrival, <i>n</i> (%)	609 (57)	1690 (55)	382 (54)	200 (53)	<i>p</i> = 0.540	2881 (55)
Missing	7 (1)	14 (1)	1 (0)	11 (3)		33 (1)
AED/ EMS defibrillation, <i>n</i> (%)	501 (47)	1300 (43)	459 (65)	175 (46)	<i>p</i> < 0.001	2435 (47)
Missing	3 (0)	10 (0)	1 (0)	17 (5)		31 (1)
ROSC before transportation, <i>n</i> (%)					<i>p</i> < 0.001	

Yes	360 (34)	867 (28)	425 (60)	94 (25)		1746 (33)
No	224 (21)	700 (23)	32 (5)	41 (11)		997 (19)
N/A***	479 (45)	1474 (48)	251 (35)	217 (58)		2421 (46)
Missing	9 (1)	21 (1)	3 (0)	25 (7)		58 (1)
Result prehospital CPR, <i>n</i> (%)					<i>p</i> < 0.001	
ROSC	287 (27)	622 (20)	394 (55)	81 (22)		1384 (27)
Transport to hospital continuing CPR	290 (27)	946 (31)	87 (12)	68 (18)		1391 (27)
Deceased	486 (45)	1479 (48)	227 (32)	205 (54)		2397 (46)
Missing	9 (1)	15 (1)	3 (0)	23 (6)		50 (1)
Alert to start CPR by EMS mm:ss, Median (IQR)	10:49 (8:47–13:25)	11:25 (9:21–13:54)	11:26 (9:06–13:53)	11:24 (9:29–13:44)	<i>p</i> < 0.001	11:16 (9:14–13:47)
Missing	133 (12)	346 (11)	159 (22)	107 (28)		745 (14)
Alert to ROSC mm:ss, Median (IQR)	23:30 (18:24–28:38)	23:30 (18:36–29:10)	15:36 (12:09–22:32)	22:32 (13:52–33:59)	<i>p</i> < 0.001	22:26 (16:49–28:27)
Missing or no ROSC	712 (66)	2140 (70)	401 (56)	288 (76)		3541 (68)
Start CPR by EMS to ROSC mm:ss, Median (IQR)	12:06 (8:21–17:30)	11:56 (7:55–17:14)	4:09 (1:55–10:31)	9:03 (3:25–18:03)	<i>p</i> < 0.001	10:54 (6:03–16:45)
Missing or no ROSC	712 (66)	2140 (70)	405 (57)	291 (77)		3548 (68)

Abbreviations: AED= automated external defibrillator; BLS= basic life support; BVM= bag-valve-mask; CPR= cardiopulmonary resuscitation; EMS= emergency medical services; ETT= endotracheal tube; IQR= interquartile range; SAD= supraglottic airway device; OHCA= out-of-hospital cardiac arrest; ROSC= return of spontaneous circulation; VF= ventricular fibrillation; VT= ventricular tachycardia.

*For cases witnessed by EMS, the initial rhythm displayed on the ambulance monitor may not reflect the true first recorded rhythm during the arrest.

**For some cases, the first monitored rhythm was either missing or could not be clearly classified.

***N/A (not applicable): applies to patients where ROSC prior to transport could not occur, such as EMS-witnessed arrests or cases where EMS determined that transport was unnecessary.

Patients for whom BVM was maintained as the final airway device were generally younger than those who received advanced airway management, with median ages of 67 versus 71 years. This group also had a higher proportion of EMS-witnessed arrests (21% compared with 7%), more frequently presented with a shockable initial rhythm (42% versus 29%), and achieved return of spontaneous circulation (ROSC) more often (55% versus 22%; **Table 1**). The median duration from the start of CPR by EMS to ROSC was shorter in the BVM group at 4 minutes 9 seconds (IQR 1:55–10:31) versus 11 minutes 56 seconds (IQR 8:03–17:19) for patients managed with an advanced airway. Among those whose resuscitation ended with BVM, an attempt to insert an advanced airway occurred in 120 patients (17%), comprising 13% SAD attempts and 11% ETT attempts, whereas no attempt was made in 568 patients (80%). In 23 cases (3%), data on airway attempts were not recorded (**Table 2**).

Table 2. Airway characteristics of included cases stratified by definitive airway device.

	ETT	BVM	Unknown	SAD	Statistics	Total
Adult OHCA with CPR attempt, <i>n</i> (%; 95% CI)	1072 (21; 19–22)	711 (14; 13–15)	377 (7; 7–8)	3062 (59; 57–60)		5222
SAD attempted, <i>n</i> (%)	112 (10)	90 (13)	7 (2)	3062 (100)	<i>p</i> < 0.001	3271 (63)
Missing	40 (4)	23 (3)	365 (97)	0		428 (8)
First pass success SAD, <i>n</i> (% of attempted)	44 (39)	1 (1)	4 (57)	3002 (98)	<i>p</i> < 0.001	3051 (93)
Tube attempted, <i>n</i> (%)	1072 (100)	75 (11)	11 (3)	498 (16)	<i>p</i> < 0.001	1656 (32)
Missing	0	24 (3)	358 (95)	180 (6)		562 (11)
First pass success tube, <i>n</i> (% of attempted)	1017 (95)	0 (0)	4 (36)	3 (1)	<i>p</i> < 0.001	1024 (62)
Order of successful airway devices, <i>n</i> (%)					<i>p</i> < 0.001	
BVM	0	708 (100)	0	0		708 (14)
BVM + SAD	0	0	0	1996 (65)		1996 (38)
BVM + ETT	871 (81)	0	0	0		871 (17)
BVM + SAD + ETT	39 (4)	0	0	0		39 (1)
BVM + ETT + SAD	0	0	0	2 (0)		2 (0)

SAD	0	0	0	1003 (33)	1003 (19)
SAD + ETT	5 (1)	0	0	0	5 (0)
ETT	38 (4)	0	0	0	38 (1)
ETT + SAD	0	0	0	1 (0)	1 (0)
Missing	119 (11)	3 (0)	377 (100)	60 (2)	559 (11)

BVM= bag-valve-mask, CPR= cardiopulmonary resuscitation, ETT= endotracheal tube, SAD= supraglottic airway device, OHCA= out-of-hospital cardiac arrest.

Patients receiving an SAD had lower occurrences of witnessed arrest (65 percent vs. 70 percent), AED or EMS defibrillation (43% vs. 47%), ROSC prior to transport (28% vs. 34%), and arrival at hospital with ROSC (20 percent vs. 27 percent) compared with those managed with an ETT, while ongoing CPR during transport was more frequent with an SAD (31% vs. 27%). Baseline characteristics of OHCA patients over time are provided in Supplementary Table A. First-attempt success for SAD insertion was high at 93% and remained stable across the years (**Table 3, Figure 2A**). In 44 instances (representing 1% of successful SAD cases), intubation was subsequently performed and succeeded. Overall, first-pass intubation success averaged 62%, showing a trend toward improvement over the years (from 53% in 2019 to 68 percent in 2023), even as the proportion of cases where intubation was attempted declined (61% in 2019 to 29 percent in 2023; **Figure 2B**).

Table 3. Airway characteristics of included cases stratified by year of arrest.

	2023	2022	2021	2020	2019	Statistics	Total
Adult OHCA with CPR attempt, <i>n</i> (%)	1007 (19)	1093 (21)	1111 (21)	1000 (19)	1011 (19)		5222
SAD attempted, <i>n</i> (%)	627 (62)	685 (63)	801 (72)	694 (69)	463 (46)	$p < 0.001$	3270 (63)
Missing	84 (8)	86 (8)	95 (9)	61 (6)	103 (10)		429 (8)
First pass success SAD, <i>n</i> (%) of attempted)	578 (92)	647 (95)	740 (92)	650 (94)	436 (94)	$p = 0.378$ (X20.378 (X	3051 (93)
Tube attempted, <i>n</i> (%)	291 (29)	288 (26)	211 (19)	249 (25)	617 (61)	$p < 0.001$	1656 (32)
Missing	80 (8)	90 (8)	99 (9)	107 (11)	186 (18)		562 (11)
First pass success tube, <i>n</i> (%) of attempted)	197 (68)	197 (68)	144 (68)	161 (65)	325 (53)	$p < 0.001$	1024 (62)
Definitive device, <i>n</i> (%; 95% CI)						$p < 0.001$	
BVM	132 (13; 11–15)	158 (15; 13–17)	121 (11; 9–13)	131 (13; 11–15)	169 (17; 15–19)		711 (14; 13–15)
SAD	581 (58; 55–61)	644 (59; 56–62)	752 (68; 65–70)	651 (65; 62–68)	434 (43; 40–46)		3062 (59; 57–60)
ETT	209 (21; 18–23)	211 (19; 17–22)	156 (14; 12–16)	167 (17; 15–19)	329 (33; 30–36)		1072 (21; 19–22)
Unknown (missing)	85 (8; 7–10)	80 (7; 6–9)	82 (7; 6–9)	51 (5; 4–7)	79 (8; 6–10)		377 (7; 7–8)
Order of successful airway devices, <i>n</i> (%)						$p = 0.005$	
BVM	130 (13)	157 (14)	121 (11)	131 (13)	169 (17)		708 (14)
BVM + SAD	533 (53)	530 (49)	244 (22)	257 (26)	432 (43)		1996 (38)
BVM + ETT	195 (19)	161 (15)	71 (6)	119 (12)	325 (32)		871 (17)
BVM + SAD + ETT	9 (1)	15 (1)	5 (1)	7 (1)	3 (0)		39 (1)
BVM + ETT + SAD	0 (0)	1 (0)	1 (0)	0 (0)	0 (0)		2 (0)
SAD	46 (5)	99 (9)	488 (44)	370 (37)	0 (0)		1003 (19)
SAD + ETT	1 (0)	0 (0)	1 (0)	3 (0)	0 (0)		5 (0)
ETT	1 (0)	12 (1)	10 (1)	15 (2)	0 (0)		38 (1)
ETT + SAD	0 (0)	0 (0)	1 (0)	0 (0)	0 (0)		1 (0)
Missing	92 (9)	118 (11)	169 (15)	98 (10)	82 (8)		559 (11)

BVM= bag-valve-mask, CPR= cardiopulmonary resuscitation, ETT= endotracheal tube, SAD= supraglottic airway device, OHCA= out-of-hospital cardiac arrest.




Figure 2. Trends in airway management over the years: (A) shows the proportion of patients in whom an SAD was attempted and successfully inserted on the first pass, and (B) shows the proportion of patients in whom an ETT was attempted and successfully placed on the first pass. ETT refers to endotracheal tube, and SAD to supraglottic airway device. Among OHCA patients, the most frequently successful airway approach was BVM ventilation, followed by SAD placement (38 percent). During the period of airway-related COVID restrictions, direct SAD placement partially replaced strategies that initially relied on BVM ventilation, and this approach continued even after the restrictions were lifted. Direct ETT insertion as an initial advanced airway strategy was not commonly used during the COVID period. The sequence of successful airway management approaches is summarized in a Sankey diagram (Figure 3). In one case from the BVM group, an SAD was successfully placed, followed by an unsuccessful intubation attempt, after which BVM was resumed without further advanced airway attempts. Among patients whose first advanced airway was a successfully placed ETT, three later received a successfully placed SAD; in one case, the switch was likely due to esophageal intubation indicated by persistently low end-tidal CO₂, while no further details were available for the remaining two cases




Figure 3. Sankey diagram showing the sequence of successful airway management, with BVM indicating bag-valve-mask, ETT for endotracheal tube, and SAD for supraglottic airway device

Between 2019 and 2023, the use of ETT as the definitive airway decreased from 33 percent (95 percent CI: 30–36 percent) to 21 percent (95 percent CI: 18–23 percent), whereas SAD usage increased from 43 percent (95 percent CI: 40–46 percent) to 58 percent (95 percent CI: 55–61 percent). Over the same period, ETT first-pass success improved from 53 percent in 2019 to 68 percent in 2023, despite fewer intubation attempts, while SAD first-pass success remained consistently high at an average of 93%. Similar findings during the COVID-19 pandemic, reported by Armour *et al.*, showed that reduced ETT attempts did not negatively impact intubation success [40]. COVID-19 restrictions led to decreased ETT use and increased SAD use, as also observed in a meta-analysis by Kim *et al.* evaluating prehospital airway management during the pandemic [41], and this pattern persisted even after returning to pre-pandemic guidelines. In this study, the enforced use of primary SAD placement may have allowed EMS nurses to gain more experience and shifted preference toward SAD as the initial airway device. Post-pandemic, selection may have occurred both in patient complexity and among EMS providers with higher intubation skills, contributing to the observed improvement in ETT first-pass success. Although the current ambulance protocol primarily recommends SAD insertion before ETT, only 10 percent of patients ultimately managed with ETT underwent an initial SAD attempt. While the proportion of patients receiving an SAD attempt increased from 46 percent in 2019 to 62 percent in 2023, at least 29 percent of patients

in 2023 did not have an SAD attempt even after the protocol update in May 2023. The rationale for airway device selection was not recorded; EMS providers may have deemed SAD unsuitable due to factors such as aspiration, which occurs in up to 32% of OHCA cases, with oral fluids often prompting deviation from the SAD-first approach as reported in the AIRWAYS-2 trial [4,13,42–45]. Paramedics are roughly five times more likely to use an ETT than SAD in cases of regurgitation. However, the incidence of regurgitation and aspiration does not significantly differ between SAD and ETT during OHCA [4], supporting the notion that primary SAD attempts remain a reasonable strategy. Previous research by Berdowski demonstrated that full implementation of national resuscitation guidelines can take up to 1.5 years, suggesting potential improvements in adherence over time [46]. This study has several limitations. First, airway management strategies and success rates were self-reported by ambulance nurses, introducing potential bias, likely inflating first-pass success. Variability in the definition of first-pass success may have further contributed to this bias [47]. Second, the study describes EMS airway management processes without evaluating the reasons behind device selection or patient outcomes, limiting conclusions regarding the appropriateness of airway choices. Nonetheless, the findings may guide targeted airway management training or development of new strategies and devices.

Although the results support primary SAD attempts for EMS, intubation remains necessary for certain patients, including those with failed SAD placement (~3%) or those who required ETT after initial SAD success (1%). Improving ETT first-pass success is critical, as over one-third of patients were not intubated successfully on the first attempt. First-pass ETT success in OHCA ranges from 51 percent to 90 percent, depending on definition, setting, and provider training [3, 4, 13, 23, 48–50]. The current cohort's success falls within this range but is not considered "high" per 2021 ERC guidelines. Ongoing intubation training is essential, and adjuncts such as bougies or videolaryngoscopy should be considered [51, 52]. Videolaryngoscopy has been shown to enhance first-pass success for less experienced providers and is associated with improved survival [53–57].

Future randomized trials should examine the impact of videolaryngoscopy versus direct laryngoscopy on first-pass success and outcomes. Additionally, comparing primary SAD versus BVM-first strategies on ventilation effectiveness and outcomes warrants further investigation, as ventilation quality is the ultimate goal of airway management, not the device itself [58, 59].

Conclusion

In recent years, the primary advanced airway device in OHCA shifted from ETT to SAD. First-pass success for SADs remained high, and few patients required intubation after initial SAD placement. Interestingly, ETT first-pass success improved despite fewer intubation attempts. These findings support continued use of primary SAD strategies by EMS, though intubation remains essential for a subset of patients, highlighting the need for strategies to improve ETT first-pass success.

Acknowledgments: We would like to thank all datamanagers at ARREST for their careful handling of the data, and the ambulance nurses, EMS Medical Directors, police and fire department, and HartslagNu for collaborating with ARREST.

Conflict of interest: The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: 'LD and HvS report a grant to their institution from Stryker Emergency Care. JdJ and VE report no conflict of interest. PS reports a grant from Health Holland, outside the scope of this study. MWH reports grants to his institution from ZonMW and ESAIC, and consulting fees paid to his institution from IDD Pharma, Medical Developments and PAION, all outside the scope of this study.'

Financial support: The ARREST registry is maintained by an unconditional grant of Stryker Emergency Care, Redmond, WA. This study has not received specific funding.

Ethics statement: None.

References

1. Lurie KG, Nemergut EC, Yannopoulos D, Sweeney M. The physiology of cardiopulmonary resuscitation. *Anesth Analg*. 2016;122:767–83. doi:10.1213/ANE.0000000000000926
2. van Schuppen H, Boomars R, Kooij FO, den Tex P, Koster RW, Hollmann MW. Optimizing airway management and ventilation during prehospital advanced life support in out-of-hospital cardiac arrest: a narrative review. *Best Pract Res Clin Anaesthesiol*. 2021;35:67–82. doi:10.1016/j.bpa.2020.11.003
3. Wang CH, Lee AF, Chang WT, Chen WJ, Chien CY, Lin JW, et al. Comparing effectiveness of initial airway interventions for out-of-hospital cardiac arrest: a systematic review and network meta-analysis of clinical controlled trials. *Ann Emerg Med*. 2020;75:627–36. doi:10.1016/j.annemergmed.2019.12.003

4. Bengier JR, Kirby K, Black S, Perkins GD, Brett SJ, Clout M, et al. Effect of a strategy of a supraglottic airway device vs tracheal intubation during out-of-hospital cardiac arrest on functional outcome: the AIRWAYS-2 randomized clinical trial. *JAMA*. 2018;320:779–91. doi:10.1001/jama.2018.11597
5. Lesnick JA, Moore JX, Zhang Y, Patterson T, Ferguson L, Shults J, et al. Airway insertion first pass success and patient outcomes in adult out-of-hospital cardiac arrest: the pragmatic airway resuscitation trial. *Resuscitation*. 2021;158:151–6. doi:10.1016/j.resuscitation.2020.11.030
6. Wang HE, Simeone SJ, Weaver MD, Callaway CW. Interruptions in cardiopulmonary resuscitation from paramedic endotracheal intubation. *Ann Emerg Med*. 2009;54:645–652.e1. doi:10.1016/j.annemergmed.2009.05.024
7. Murphy DL, King JA, Blackwood J, van Haren F, Donnelly JP, Ward MJ, et al. The apnea interval: ventilation interruption during tracheal intubation and its association with cardiac arrest resuscitation care and outcome. *Resuscitation*. 2025;176:110588. doi:10.1016/j.resuscitation.2025.110588
8. Nakanishi T, Sakamoto S, Yoshimura M, Fujiwara K, Toriumi T. Learning curve of i-gel insertion in novices using a cumulative sum analysis. *Sci Rep*. 2023;13:7121. doi:10.1038/s41598-023-34152-5
9. Konrad C, Schupfer G, Wietlisbach M, Gerber H. Learning manual skills in anesthesiology: is there a recommended number of cases for anesthetic procedures? *Anesth Analg*. 1998;86:635–9. doi:10.1097/0000539-199803000-00037
10. de Oliveira Filho GR. The construction of learning curves for basic skills in anesthetic procedures: an application for the cumulative sum method. *Anesth Analg*. 2002;95:411–416. doi:10.1097/0000539-200208000-00033
11. Komatsu R, Kasuya Y, Yogo H, Sugimoto T, Yamakage M, Hashimoto H, et al. Learning curves for bag-and-mask ventilation and orotracheal intubation: an application of the cumulative sum method. *Anesthesiology*. 2010;112:1525–31. doi:10.1097/ALN.0b013e3181d96779
12. Bengier JR, Lazaroo MJ, Clout M, Perkins GD, Brett SJ, Black S, et al. Randomized trial of the i-gel supraglottic airway device versus tracheal intubation during out of hospital cardiac arrest (AIRWAYS-2): patient outcomes at three and six months. *Resuscitation*. 2020;157:74–82. doi:10.1016/j.resuscitation.2020.09.026
13. Wang HE, Schmicker RH, Daya MR, Rea TD, Berg RA, Hostler D, et al. Effect of a strategy of initial laryngeal tube insertion vs endotracheal intubation on 72-hour survival in adults with out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA*. 2018;320:769–78. doi:10.1001/jama.2018.7044
14. Lee AF, Chien YC, Lee BC, Chen KH, Chang WT, Chien CY, et al. Effect of placement of a supraglottic airway device vs endotracheal intubation on return of spontaneous circulation in adults with out-of-hospital cardiac arrest in Taipei, Taiwan: a cluster randomized clinical trial. *JAMA Netw Open*. 2022;5:e2148871. doi:10.1001/jamanetworkopen.2021.48871
15. in 't Veld C, van Exter P, Rombouts M, de Visser M, de Vos R. Landelijk protocol ambulancezorg 8 [Internet]. 8 ed. Zwolle: Ambulancezorg Nederland; 2014. [cited 2025 May 30]. Available from: Visio-20140716 LPA8 boek.vsd
16. in 't Veld C, van Exter P, Rombouts M, de Visser M, de Vos R. Landelijk protocol ambulancezorg 8.1 [Internet]. 8.1 ed. Zwolle: Ambulancezorg Nederland; 2016. [cited 2025 May 30]. Available from: https://www.ambulancezorg.nl/static/upload/raw/d24e66de-8f05-47a1-92cd-3ec0278c2287/Ambulancezorg_Nederland_LPA8.1_herdruk_2019.oktober.wm.bv.pdf
17. Brintjes A, Caminanda K, Dercksen B, Van Duin T, Lelieveld K, Ruteen M, et al. Landelijk protocol ambulancezorg 9. 9 ed. Zwolle: Ambulancezorg Nederland; 2023. [cited 2025 May 30]. Available from: <https://www.ambulancezorg.nl/static/upload/raw/cd6c501a-9ed7-4b4f-ad02-570813341e41/Onderwijsboek+protocollen+LPA9+2023.TLP-Wit.pdf>
18. Cady CE, Pirrallo RG. The effect of combitube use on paramedic experience in endotracheal intubation. *Am J Emerg Med*. 2005;23:868–71. doi:10.1016/j.ajem.2005.07.013
19. Stewart RD, Paris PM, Pelton GH, Garretson D. Effect of varied training techniques on field endotracheal intubation success rates. *Ann Emerg Med*. 1984;13:1032–6. doi:10.1016/s0196-0644(84)80064-5
20. Warner KJ, Carlbom D, Cooke CR, Bulger EM, Copass MK, Sharar SR. Paramedic training for proficient prehospital endotracheal intubation. *Prehosp Emerg Care*. 2010;14:103–8. doi:10.3109/10903120903144858
21. Garza AG, Gratton MC, Coontz D, Noble E, Ma OJ. Effect of paramedic experience on orotracheal intubation success rates. *J Emerg Med*. 2003;25:251–6. doi:10.1016/s0736-4679(03)00198-7
22. Doran JV, Tortella BJ, Drivet WJ, Lavery RF. Factors influencing successful intubation in the prehospital setting. *Prehosp Disaster Med*. 1995;10:259–64. doi:10.1017/s1049023x00042138
23. Dyson K, Bray J, Smith K, Bernard S, Finn J. A systematic review of the effect of emergency medical service practitioners' experience and exposure to out-of-hospital cardiac arrest on patient survival and procedural performance. *Resuscitation*. 2014;85:1134–41. doi:10.1016/j.resuscitation.2014.05.020

24. Bierowski AE, Comber PC, Kuc A, Shah A, Carroll G. The effect of prehospital protocol modification during COVID-19 on first-pass intubation success rates. *Prehosp Disaster Med.* 2025;1–4. doi:10.1017/S1049023X25000238
25. Thomas J, Crowe R, Schulz K, Nehme Z, Smith K, Bray J, et al. Association between emergency medical service agency intubation rate and intubation success. *Ann Emerg Med.* 2024;84:1–8. doi:10.1016/j.annemergmed.2023.11.005
26. Gage CB, Powell JR, Nassal M, Wang H, Panchal AR. Secular trends in airway management of out-of-hospital cardiac arrest in the National Emergency Medical Services Information System (NEMSIS) dataset. *Resuscitation.* 2023;193:110024. doi:10.1016/j.resuscitation.2023.110024
27. Wang HE, Yu MI, Crowe RP, Schmicker RH, Daya MR, Rea TD, et al. Longitudinal changes in emergency medical services advanced airway management. *JAMA Netw Open.* 2024;7:e27763. doi:10.1001/jamanetworkopen.2024.27763
28. Stieglis R, Zijlstra JA, Riedijk F, Oving I, van der Wal G, Tijssen JG, et al. Alert system-supported lay defibrillation and basic life-support for cardiac arrest at home. *Eur Heart J.* 2022;43:1465–74. doi:10.1093/eurheartj/ehab802
29. Grasner JT, Bray JE, Nolan JP, Berdowski J, Herlitz J, Wingen S, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: 2024 update of the Utstein out-of-hospital cardiac arrest registry template. *Resuscitation.* 2024;201:110288. doi:10.1016/j.resuscitation.2024.110288
30. Christiaans H, van Eenennaam F, Hoogeveen M, et al. MMT Inzet- en cancelcriteria [Internet]. Zeist: Landelijk netwerk acute zorg, Ambulancezorg Nederland; 2013. [cited 2025 May 30]. Available from: [Inzet- en cancelcriteria_MMT-_LNAZ-AZN.PDF](#)
31. Ali S, Moors X, van Schuppen H, Mommers L, Weelink E, Meuwese CL, et al. A national multi centre pre-hospital ECPR stepped wedge study; design and rationale of the ON-SCENE study. *Scand J Trauma Resusc Emerg Med.* 2024;32:31. doi:10.1186/s13049-024-01198-x
32. Soar J, Böttiger BW, Carli P, Couper K, Deakin CD, Djävrv T, et al. European Resuscitation Council guidelines 2021: adult advanced life support. *Resuscitation.* 2021;161:115–51. doi:10.1016/j.resuscitation.2021.02.010
33. NRR. Reanimatie buiten het ziekenhuis tijdens de COVID-19 pandemie. Advies voor tijdelijke aanpassingen reanimatie richtlijnen in verband met de COVID-19 pandemie [Internet]. Voorschoten: Nederlandse Reanimatie Raad; 2020. [cited 2025 May 30]. Available from: [ADVIES-NRR-OHCA_COVID19_versie2.7_200407.pdf](#)
34. NRR. Reanimatie buiten het ziekenhuis tijdens de COVID-19 pandemie. Herziend advies voor tijdelijke aanpassingen reanimatie richtlijnen in verband met de COVID-19 pandemie [Internet]. Voorschoten: Nederlandse Reanimatie Raad; 2020. [cited 2025 May 30]. Available from: https://www.ambulancezorg.nl/static/upload/raw/9cfce8df-601f-48ee-8025-92dbc3a39b70/Herziend-ADVIES-NRR-OHCA_COVID19_versie+19+juni+2020.pdf
35. NRR. Reanimatie buiten het ziekenhuis tijdens de COVID-19 pandemie. Advies voor tijdelijke aanpassingen reanimatie richtlijnen in verband met de COVID-19 pandemie [Internet]. Voorschoten: Nederlandse Reanimatie Raad; 2021. [cited 2025 May 30]. Available from: https://www.reanimatieraad.nl/app/uploads/2021/11/Advies_OHCA-covid19pandemie_23nov2021.pdf
36. Blom MT, van Hoeijen DA, Bardai A, Berdowski J, Souverein PC, De Bruin ML, et al. Genetic, clinical and pharmacological determinants of out-of-hospital cardiac arrest: rationale and outline of the AmsterDdam Resuscitation Studies (ARREST) registry. *Open Heart.* 2014;1:e000112. Available from: <http://dx.doi.org/10.1136/openhrt-2014-000112>. doi:10.1136/openhrt-2014-000112
37. Centraal Bureau voor de Statistiek. Inwonertal per provincie 1 september 2022 [Internet]. 2023 Jul [cited 2025 May 30]. Available from: <https://www.cbs.nl/nl-nl/maatwerk/2023/07/inwonertal-per-provincie-1-september-2022>
38. Perkins GD, Jacobs IG, Nadkarni VM, Berg RA, Bhanji F, Biarent D, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein Resuscitation Registry Templates for Out-of-Hospital Cardiac arrest: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia); and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Circulation.* 2015;132:1286–300. doi:10.1161/CIR.000000000000144
39. Nam JM. A simple approximation for calculating sample sizes for detecting linear trend in proportions. *Biometrics.* 1987;43:701–5.

40. Armour R, Ghamarian E, Helmer J, Buick JE, Thorpe K, Austin M, et al. Impact of the COVID-19 pandemic on Canadian emergency medical system management of out-of-hospital cardiac arrest: a retrospective cohort study. *Resuscitation*. 2024;194:110054. doi:10.1016/j.resuscitation.2023.110054
41. Kim JH, Ahn C, Park Y, Won M. Comparison of out-of-hospital cardiac arrests during the COVID-19 pandemic with those before the pandemic: an updated systematic review and meta-analysis. *Front Public Health*. 2023;11:1180511. doi:10.3389/fpubh.2023.1180511
42. Virkkunen I, Ryyänen S, Kujala S, Ala-Kokko T, Silfvast T, Pettilä V, et al. Incidence of regurgitation and pulmonary aspiration of gastric contents in survivors from out-of-hospital cardiac arrest. *Acta Anaesthesiol Scand*. 2007;51:202–205. doi:10.1111/j.1399-6576.2006.01229.x
43. Simons RW, Rea TD, Becker LJ, Eisenberg MS. The incidence and significance of emesis associated with out-of-hospital cardiac arrest. *Resuscitation*. 2007;74:427–431. doi:10.1016/j.resuscitation.2007.01.038
44. Stone BJ, Chantler PJ, Baskett PJF. The incidence of regurgitation during cardiopulmonary resuscitation: a comparison between the bag valve mask and laryngeal mask airway. *Resuscitation*. 1998;38:3–6. doi:10.1016/s0300-9572(98)00068-9
45. Kirby K, Brandling J, Robinson M, Thomas M, Voss S, Bengner J. The experiences of EMS providers taking part in a large randomised trial of airway management during out of hospital cardiac arrest, and the impact on their views and practice. Results of a survey and telephone interviews. *Resuscitation*. 2020;149:1–9. doi:10.1016/j.resuscitation.2020.01.034
46. Berdowski J, Schmohl A, Tijssen JGP, Koster RW. Time needed for a regional emergency medical system to implement resuscitation guidelines 2005—the Netherlands experience. *Resuscitation*. 2009;80:1336–1341. doi:10.1016/j.resuscitation.2009.08.011
47. Nauka PC, Moskowitz A, Fein DG. Appraising first-pass success: during emergency airway management, what does it mean to be successful? *Ann Am Thorac Soc*. 2023;20:21–23. doi:10.1513/AnnalsATS.202208-661VP
48. Goldenberg IF, Campion BC, Siebold CM, McBride JW, Long LA. Esophageal gastric tube airway vs endotracheal tube in prehospital cardiopulmonary arrest. *Chest*. 1986;90:90–96. doi:10.1378/chest.90.1.90
49. Mulder PJ, Oetomo SB, Vloet L, De Vries P, Hoogerwerf N. Comparison in effectiveness and safety between a supraglottic airway device and endotracheal intubation in out-of-hospital cardiac arrest in the Netherlands. *Resuscitation*. 2013;84:S17.
50. Levi D, Hoogendoorn J, Samuels S, Maguire L, Troncoso R, Gunn S, et al. The i-gel(®) supraglottic airway device compared to endotracheal intubation as the initial prehospital advanced airway device: a natural experiment during the COVID-19 pandemic. *J Am Coll Emerg Physicians Open*. 2024;5:e13150. doi:10.1002/emp2.13150
51. Latimer AJ, Harrington B, Counts CR, Ruark K, Maynard C, Watase T, et al. Routine use of a bougie improves first-attempt intubation success in the out-of-hospital setting. *Ann Emerg Med*. 2021;77:296–304. doi:10.1016/j.annemergmed.2020.10.016
52. Xu X, Ma H, Zhang Y, Liu W, Jung B, Li X, et al. Efficacy of bougie first approach for endotracheal intubation with video laryngoscopy during continuous chest compression: a randomized crossover manikin trial. *BMC Anesthesiol*. 2024;24:181. doi:10.1186/s12871-024-02560-3
53. Chien YT, Ong JR, Tam KW, Loh EW. Video laryngoscopy and direct laryngoscopy for cardiac arrest: a meta-analysis of clinical studies and trials. *Am J Emerg Med*. 2023;73:116–24. doi:10.1016/j.ajem.2023.08.028
54. Maissan I, van Lieshout E, de Jong T, van Vledder M, Houmes RJ, Hartog DD, et al. The impact of video laryngoscopy on the first-pass success rate of prehospital endotracheal intubation in the Netherlands: a retrospective observational study. *Eur J Trauma Emerg Surg*. 2022;48:4205–4213. doi:10.1007/s00068-022-01962-7
55. van Schuppen H, Wojciechowicz K, Hollmann MW, Preckel B. Tracheal intubation during advanced life support using direct laryngoscopy versus Glidescope(®) videolaryngoscopy by clinicians with limited intubation experience: a systematic review and meta-analysis. *J Clin Med*. 2022;11:6291. doi:10.3390/jcm11216291
56. Phillips JP, Anger DJ, Rogerson MCT, Myers LA, McCoy RG. Transitioning from direct to video laryngoscopy during the COVID-19 pandemic was associated with a higher endotracheal intubation success rate. *Prehosp Emerg Care*. 2024;28:200–8. doi:10.1080/10903127.2023.2175087
57. Risse J, Fischer M, Meggiolaro KM, Fariq-Spiegel K, Pabst D, Manegold R, et al. Effect of video laryngoscopy for non-trauma out-of-hospital cardiac arrest on clinical outcome: a registry-based analysis. *Resuscitation*. 2023;185:109688. doi:10.1016/j.resuscitation.2023.109688
58. Segond N, Wittig J, Kern WJ. Towards a common terminology of ventilation during cardiopulmonary resuscitation. *Resuscitation*. 2025;191:110511. doi:10.1016/j.resuscitation.2025.110511

59. van Eijk JA, Doeleman LC, Loer SA, Koster RW, van Schuppen H, Schober P. Ventilation during cardiopulmonary resuscitation: a narrative review. *Resuscitation*. 2024;203:110366. doi:10.1016/j.resuscitation.2024.110366