

Prevalence of Palliative Needs and Screening Accuracy of Psychological Symptoms Using the Integrated Palliative Care Outcome Scale

Pamela D Moore¹, Ayuk Patricia^{2*}, Zhe-Sheng Chen²

¹*Department of Sociology and Gerontology, Miami University, 375 Upham Hall 100 Bishop Circle, Oxford, OH, 45056, USA.*

²*Prisma Health Richland Hospital, 5 Richland Medical Park Dr., Columbia, SC, 29203, USA.*

Abstract

Individuals receiving palliative care often face substantial symptom burdens that affect both themselves and their families. Emotional distress, including depression and anxiety, should not be considered a normal part of advanced illness; it requires routine assessment, timely intervention, and ongoing monitoring. Psychological distress has been associated with worse physical symptoms, increased suffering, and higher mortality in patients with cancer. Comprehensive yet concise tools are needed to capture both physical and psychosocial needs, and the *Integrated Palliative care Outcome Scale* (IPOS) is one such measure. This study aimed to evaluate the palliative care needs of patients and to examine how accurately IPOS items related to anxiety and depression identify psychological distress. A multicentre observational study was conducted in nine Portuguese healthcare settings using convenience sampling. Eligible participants were adults (≥ 18 years) with a diagnosis of incurable or life-limiting illness who were cognitively able to consent. Patients experiencing acute distress or cognitive impairment were excluded. Descriptive statistics summarized participant characteristics. The diagnostic performance of IPOS anxiety and depression items was evaluated against the *Hospital Anxiety and Depression Scale* (HADS) using Receiver Operating Characteristic (ROC) curves and Area Under the Curve (AUC) analysis. Of 1,703 individuals screened between July 2015 and February 2016, 135 (7.9%) met inclusion criteria. The majority were excluded for being healthy (75.2%), largely from high-volume primary care centres. Participants had a mean age of 66.8 years (SD 12.7), with 43% female, and 80.7% had a cancer diagnosis. The most frequently reported concerns included anxiety among family or friends (36.3%), personal anxiety about illness (13.3%), and feelings of depression (9.6%). Applying a threshold score of 2/3, the IPOS items for anxiety and depression achieved AUC values above 70%, demonstrating acceptable screening accuracy. Psychological, familial, and spiritual needs were the most prominent concerns among this palliative population, highlighting that physical symptoms may be better managed than non-physical issues. Routine implementation of the IPOS could support healthcare teams in identifying and monitoring psychosocial distress, improving holistic, patient-centred care for both patients and their families.

Keywords: Palliative care, Psychological distress, Depression, Anxiety, Screening accuracy, Patient-reported outcomes

Introduction

Palliative care adopts a holistic approach that can be introduced early alongside disease-directed treatments [1], aiming to alleviate both physical and non-physical symptoms for patients and their families [2–7]. While physical symptoms are often easier for healthcare professionals, patients, and caregivers to identify, non-physical symptoms can equally impair quality of life and contribute to suffering [8]. Evidence from a rapid systematic

Corresponding author: Ayuk Patricia

Address: Prisma Health Richland Hospital, 5 Richland Medical Park Dr., Columbia, SC, 29203, USA.

E-mail: ✉ Ayukpatricia1980@yahoo.com

Received: 14 September 2021; **Revised:** 29 October 2021;

Accepted: 05 December 2021; **Published:** 22 December 2021

How to Cite This Article: Moore PD, Patricia A, Chen ZS. Prevalence of Palliative Needs and Screening Accuracy of Psychological Symptoms Using the Integrated Palliative Care Outcome Scale. *J Integr Nurs Palliat Care*. 2021;2:66-74. <https://doi.org/10.51847/WBgNCX5SZy>

review indicates that psychological interventions in patients with long-term conditions can significantly improve at least one quality of life outcome, with effects maintained at follow-up [9]. Similarly, Ann-Yi S and colleagues reported that 24% of palliative inpatients and 19% of outpatients at a major cancer centre benefited from psychological services [8]. Psychological distress has also been associated with more severe physical symptoms, increased suffering, and higher mortality among cancer patients [10, 11]. Consequently, these needs should be assessed using validated outcome measures, and interventions tailored accordingly [12, 13].

Depression and psychological distress exemplify needs that should not be normalized in advanced disease [14]. Rather, they require systematic screening, diagnosis, and follow-up through appropriate pharmacologic, psychological, psychiatric, or combined interventions [14]. Patient-reported outcome measures are the preferred tools for assessing subjective symptoms, as patients are best positioned to describe their experiences. In cases of cognitive impairment, proxy-reported versions may be employed [15–19]. These instruments are generally brief yet multidimensional, with certain items usable to screen for common palliative needs [20].

The present study aimed to identify the primary palliative care needs of patients receiving care in Portuguese healthcare settings and to evaluate the screening performance of two IPOS items related to psychological distress [21, 22]. We hypothesized that: 1) anxiety and depression items would score highest among non-physical symptoms, and 2) the Area Under the Curve (AUC) for IPOS Item 3 (anxiety) and Item 5 (depression) would exceed 0.7 when compared to the Portuguese *Hospital Anxiety and Depression Scale* (HADS) subscale scores [23].

Materials and Methods

Patients and settings

This multicentre observational study collected data from nine Portuguese healthcare centres spanning northern to southern regions and urban to rural locations to enhance generalizability, using convenience sampling. Participating centres included seven hospital-based palliative care services, one oncology service, and one primary care facility. All patients attending these services were screened for eligibility by local healthcare professionals. Inclusion criteria were: age ≥ 18 years, ability to provide informed consent as assessed by the clinician, diagnosis of an incurable, potentially life-threatening illness, and fluency in Portuguese. Exclusion criteria included patients experiencing distress that prevented sustained conversation or those with cognitive impairment, as judged by the clinician. A standard operating procedures manual was distributed to all centres, with a local facilitator/champion overseeing adherence. The full study protocol has been published elsewhere [19].

Measures

The study used the patient-reported Portuguese version of the IPOS, which had been culturally adapted and validated for European Portuguese [21, 22]. The adaptation process involved two independent translators (clinical and non-clinical) producing initial versions, which were reconciled into a consensus version by two independent reviewers blinded to the original IPOS. This version underwent back-translation by two additional independent translators and subsequent reconciliation. Three clinical reviews were performed by a specialist palliative care physician, a specialist nurse, and a non-clinical researcher—resulting in the final Portuguese version. Discrepancies in verb tense, word choice, and response category phrasing were resolved through discussion during translation and clinical review stages.

Measures

A Portuguese adaptation of the Integrated Palliative care Outcome Scale (IPOS) was developed and evaluated for psychometric properties. Internal consistency, excluding open-ended items, showed Cronbach's alpha values ranging from 0.68 to 0.72. Inter-rater reliability between patients and healthcare professionals was assessed using intraclass correlation coefficients (ICC), which were highest for mobility (ICC = 0.726) and lowest for practical problems (ICC = 0.088). Construct validity was supported through convergent and divergent validity analyses, with Spearman's rho ranging from 0.390 to 0.631 ($p \leq 0.000$). The instrument also demonstrated sensitivity to change, as Wilcoxon signed-rank tests revealed significant differences in three symptoms between T1 and T2.

The IPOS is a concise, 19-item, multidimensional tool capturing core concerns in palliative care. Item 1 is an open-ended question asking respondents to identify their three main problems or worries in the past week (these free-text data are not reported here). Items 2–9 employ a 5-point Likert scale (0 = not at all, 1 = slightly, 2 = moderately, 3 = severely, 4 = overwhelmingly). Item 2 lists ten common physical symptoms, with the option to add up to three additional symptoms. Item 3 addresses anxiety, Item 4 asks about family or friends' worry, and Item 5 pertains to depression. Item 6 relates to feelings of peace, Item 7 to sharing emotions with significant others, Item 8 to information needs, and Item 9 to practical problems arising from illness. The patient version includes an additional question asking whether the questionnaire was completed independently or with assistance. A footnote at the end encourages patients to discuss any concerns with their healthcare professional, enhancing the tool's real-time clinical utility.

The Portuguese version of the Hospital Anxiety and Depression Scale (HADS) is a 14-item screening tool, divided into anxiety and depression subscales with seven items each, using a 4-point Likert response format. A cut-off score of 11 is recommended for both subscales. The instrument has been validated as reliable for assessing anxiety and depression across diverse medical populations [23].

Analysis

Data quality was first examined, and Little's MCAR test confirmed that missing data were missing completely at random. Descriptive statistics were used to summarize demographic and clinical characteristics. All IPOS psychological, emotional, and spiritual items (Items 3–7) were compared to the Portuguese HADS. Receiver operating characteristic (ROC) curves were generated to determine the two IPOS items with the highest discriminant ability, and a cut-off score of 2/3 was selected after preliminary evaluation. Sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), false positive rate (FPR), false negative rate (FNR), and positive and negative likelihood ratios weighted by prevalence were calculated. Thresholds of $\geq 70\%$ were considered acceptable and $\geq 80\%$ high for sensitivity, specificity, PPV, and NPV; FPR and FNR $\leq 30\%$ were considered low. All analyses used 95% confidence intervals. No formal sample size calculation was performed, as prior literature using the Palliative care Outcome Scale and HADS involved secondary analyses of multiple datasets rather than IPOS-specific evaluation.

Ethics

Ethical approval was obtained from all relevant research ethics committees, in accordance with the 1964 Helsinki Declaration and its later amendments. All participants provided written informed consent. Data were analyzed using SPSS version 22 (IBM Corp., Armonk, NY).

Results and Discussion

Between July 1, 2015, and February 2016, 1703 individuals were screened, predominantly at a primary healthcare facility. Of these, 18 patients (1.1%) declined participation, and 140 (8.2%) were excluded. A total of 135 patients (7.9%) met inclusion criteria and were enrolled in the study (**Table 1**).

Table 1. Patients screened and included in the study by participating centre

Participating Service	Number Screened (N, % of Total Screened)	Number Included (N, % of Total Included)
Oncology Hospital Service	78 (4.6%)	25 (18.5%)
Palliative Care Service 1 (North)	96 (5.6%)	9 (6.7%)
Palliative Care Service 2 (North)	28 (1.6%)	17 (12.6%)
Palliative Care Service 3 (South)	18 (1.1%)	1 (0.7%)
Palliative Care Service 4 (North)	77 (4.5%)	24 (17.8%)
Palliative Care Service 5 (South)	64 (3.8%)	17 (12.6%)
Primary Care Centre 1 (South)	119 (7.0%)	3 (2.2%)
Primary Care Centre 2 (South)	1177 (69.1%)	25 (18.5%)
Palliative Care Service 6 (Central)	46 (2.7%)	14 (10.4%)
Total	1703 (100%)	135 (100%)

The predominant reason for exclusion was that patients were considered healthy, accounting for 75.2% of screened individuals. This outcome was anticipated, as the primary care centre screened the majority of participants, reflecting its high daily patient volume, most of whom were healthy. Among the included sample ($n = 135$), the mean age was 66.8 years ($SD = 12.7$), with 58 patients (43%) being female. Educational attainment was low for many participants, with 74 (54.8%) having up to four years of formal education, and 74 (54.8%) resided in the Northern region of Portugal. The majority of patients ($n = 109$; 80.7%) had a cancer diagnosis and were recruited primarily from the seven hospital-based palliative care services (**Table 2**).

Table 2. Demographic and clinical information of participants Demographic and clinical information N(%)

Category	Subcategory	Value
Mean Age (years, SD)		66.8 (12.7)
Gender	Male	77 (57%)
Education (Years)	Reads and writes	5 (3.7%)

	4 years	81 (60%)
	6 years	20 (14.8%)
	9 years	10 (7.4%)
	10 years to college	19 (14%)
Geographical Region		
	North	74 (54.8%)
	Central	25 (18.5%)
	South	36 (26.7%)
Area		
	Urban	94 (69.6%)
	Peri-urban	31 (23%)
	Rural	10 (7.4%)
Place of Care		
	Primary care	28 (20.7%)
	Hospital services	25 (18.5%)
	Palliative care services	82 (60.7%)
Cancer Diagnosis		109 (80.7%)
Phase of Illness		
	Stable	64 (47.4%)
	Unstable	28 (20.7%)
	Deteriorating	43 (31.9%)
	Terminal	0 (0%)
Surprise Question (Life Expectancy)		
	> 1 year	37 (27.4%)
	6 months to 1 year	45 (33.3%)
	< 6 months	48 (35.6%)

Table 3 outlines the main reasons for participant ineligibility and exclusion. Most patients (n = 98, 72.6%) were invited to participate during external consultations. Of those included, nearly one-third (31.1%) completed the questionnaires independently, without external assistance.

69

Table 3. Reasons for ineligibility and exclusion

Category	Reason for Ineligibility/Exclusion	Number (N, %)
Ineligible		
	Under 18 years	91 (5.9%)
	Does not understand Portuguese	1 (0.06%)
	Cannot read or write	78 (5.0%)
	Illness with potential for cure	3 (0.2%)
	Healthy	1165 (75.2%)
	No reason provided	72 (4.6%)
Excluded		
	Distress	52 (3.4%)
	Cognitive deterioration	73 (4.7%)
	No reason provided	15 (0.9%)
Total		1550 (100%)

Analysis of missing data indicated that values were missing at random (Little's MCAR test: Chi-Square = 2452.946, DF = 2398, $p = .213$). Missingness across items ranged from 1 to 5%, which is considered manageable, with rates below 1% regarded as trivial and those exceeding 15% potentially compromising interpretability [24]. Given the non-parametric distribution typical of palliative care populations, missing values were addressed using median imputation.

Regarding the prevalence of patient needs, the IPOS items with the highest proportion of severe concern (=4) were: family or friends being anxious or worried (36.3%), feeling anxious about illness (13.3%), feeling depressed (9.6%), feeling at peace (9.6%), sharing feelings (8.9%), and pain (7.4%). Conversely, items with the lowest prevalence (=0) included vomiting (77%), shortness of breath (67.4%), nausea (65%), information needs (60.7%), practical problems (45.2%), and constipation (43%) (**Figure 1**).




Figure 1. IPOS scores for prevalence of main palliative needs

Screening for anxiety and depression

Items 3 (anxiety) and 5 (depression) demonstrated the highest areas under the curve (AUC) (**Figure 2**). The prevalence of depression was 24.4% (95% CI: 17.6–32.7%), with an AUC of 0.72 (95% CI: 0.62–0.81, $p < 0.001$) (**Figure 3**). For this item, sensitivity was 51.5% and specificity was 78.4%, with a positive predictive value (PPV) of 43.6% and a negative predictive value (NPV) of 83.3%. Regarding anxiety, the prevalence was 23.7% (95% CI: 16.9–31.9%), and the AUC was 0.70 (95% CI: 0.60–0.80, $p < 0.001$) (**Figure 4**). Sensitivity and specificity were 65.6% and 68.0%, respectively, while PPV and NPV were 38.8% and 86.4% (**Table 4**).




Figure 2. Area Under the Curve for IPOS items 3 to 7




Figure 3. Area Under the Curve for Portuguese IPOS item 5, depression




Figure 4. Area Under the Curve for Portuguese IPOS item 3, anxiety

Table 4. Estimated Values for items 3 and 5, cut-off 2/3, against the HADS subscales, using cut-off 10/11

	Sens (%)	CI _s (%)	Spec (%)	CI _s (%)	PPV (%)	CI _s (%)	NPV (%)	CI _s (%)	FPR (%)	CI _s (%)	FNR (%)	CI _s (%)	Ratios +	CI _s (%)	Ratios -	CI _s (%)
Item 3 – anxiety Prevalence: 23.7% (C.I. 16.9–31.9%)	65.6	0.47–0.81	68.0	0.58–0.77	38.8	0.26–0.53	86.4	0.77–0.93	0.61	0.47–0.74	0.14	0.07–0.23	0.64	0.43–0.95	0.16	0.09–0.27
Item 5 – depression Prevalence: 24.4% (C.I. 17.6–32.7%)	51.5	0.34–0.69	78.4	0.69–0.86	43.6	0.28–0.60	83.3	0.74–0.90	0.56	0.40–0.72	0.17	0.10–0.26	0.78	0.49–1.21	0.2	0.13–0.31

In this study, the predominant palliative care needs identified among patients were psychological, family-related, and spiritual. The Integrated Palliative care Outcome Scale (IPOS) proved effective in systematically capturing these needs. Notably, clinical teams appeared to address physical symptoms adequately, which is encouraging. Evidence suggests that once physical needs are managed, non-physical concerns become more salient and should be systematically assessed and addressed using patient-centered outcome measures [25, 26].

Among non-physical concerns, the most frequently reported issue was anxiety experienced by family or friends (36.3%), reflecting the central role of family in Portuguese culture and the common practice of involving family in healthcare decisions. Patient-reported anxiety and depression were the next most prevalent concerns, despite the exclusion of individuals in obvious distress. This aligns with existing literature indicating that anxiety often arises in patients with advanced disease due to diagnostic, treatment, and prognostic uncertainty [14], and that depression is also common in this population [27]. The prevalence rates observed in our study are consistent with findings from Ann-Yi S. *et al.* though that study focused exclusively on cancer patients [8]. Symptom clusters, combining physical and non-physical needs, are commonly observed in both cancer and non-cancer populations [28, 29]. The observed prevalence of depression in our cohort (24.4%) was higher than that reported in Antunes *et al.* using a single-item Palliative care Outcome Scale (17.5%, 95% CI 14.1–21.6%) [21], but lower than the 30% estimated by Hotopf *et al.* for depressive disorders in advanced disease [30].

The screening performance of IPOS items for anxiety (item 3) and depression (item 5) was acceptable, with AUC values exceeding 0.70, although the lower confidence interval limits were slightly below this threshold. Using a cut-off of 2/3, sensitivity was modest, indicating limited ability to identify true positive cases; however, specificity and negative predictive value were strong. These findings suggest that these IPOS items are particularly effective in ruling out true negative cases, an important component of screening [31], reinforcing the external validity of IPOS in this context.

A key limitation of the study was the absence of a diagnostic gold standard, such as a structured psychiatric interview based on the DSM-5, due to resource constraints. Instead, the HADS was used, which, while widely accepted and extensively validated for both clinical and research purposes, remains a screening rather than diagnostic tool.

Systematic use of the Portuguese IPOS could assist clinical teams in monitoring and addressing both physical and non-physical needs of patients and families. Specifically, items 3 and 5 appear suitable for screening patients with advanced illness, particularly to exclude those without anxiety or depression [31, 32].

Conclusion

Patient-centered outcome measures serve as valuable communication tools, providing a shared language for patients, families, healthcare professionals, institutions, and policymakers, ultimately supporting evidence-informed care. Establishing the screening properties of these tools enables not only improved clinical care but also the conduct of rigorous research. This study demonstrates that the Portuguese IPOS is effective in identifying psychological needs and can be reliably used to screen patients with advanced disease for anxiety and depression.

Acknowledgments: None

Conflict of interest: None

Financial support: None

Ethics statement: None

References

1. Oliver D. Improving patient outcomes through palliative care integration in other specialised health services: what we have learned so far and how can we improve? *Ann Palliat Med.* 2018;7(S3):S219-30. doi:10.21037/apm.2018.05.05
2. Clark D. Between hope and acceptance: the medicalisation of dying. *BMJ (Clinical research ed).* 2002;324(7342):905-7.
3. Bajwah S, Higginson IJ, Ross JR, Wells AU, Birring SS, Patel A, et al. specialist palliative care is more than drugs: a retrospective study of ILD patients. *Lung.* 2012;190(2):215-20.
4. Jocham HR, Dassen T, Widdershoven G, Halfens R. Quality of life in palliative care cancer patients: a literature review. *J Clin Nurs.* 2006;15(9):1188-95.
5. Stewart AL, Teno J, Patrick DL, Lynn J. The concept of quality of life of dying persons in the context of health care. *J Pain Symptom Manag.* 1999;17(2):93-108.

6. Kupfer JM, Bond EU. Patient satisfaction and patient-centered care: necessary but not equal. *JAMA*. 2012;308(2):139–40.
7. "WHO Definition of Palliative Care". World Health Organization. Retrieved July 10, 2018. Available from: <http://www.who.int/cancer/palliative/definition/en/>.
8. Ann-Yi S, Bruera E, Wu J, Liu DD, Agosta M, Williams JL, et al. Characteristics and outcomes of psychology referrals in palliative care department. *J Pain Symptom Manag*. 2018;56(3):344–51. doi:10.1016/j.jpainsymman.2018.05.022
9. Anderson N, Ozakinci G. Effectiveness of psychological interventions to improve quality of life in people with long-term conditions: rapid systematic review of randomised controlled trials. *BMC Psychol*. 2018;6(1):11.
10. Hamer M, Chida Y, Molloy GJ. Psychological distress and cancer mortality. *J Psychosom Res* 2009;66(3):255–8.
11. Krikorian a, Limonero JT, Roman JP, Vargas JJ, Palacio C. Predictors of suffering in advanced cancer. *Am J Hosp Palliat Care*. 2014;31(5):534–42.
12. Bausewein C, Daveson BA, Currow DC, Downing J, Deliens L, Radbruch L, et al. EAPC white paper on outcome measurement in palliative care: improving practice, attaining outcomes and delivering quality services – recommendations from the European Association for Palliative Care (EAPC) task force on outcome measurement. *Palliat Med*. 2016;30(1):6–22. doi:10.1177/0269216315589898
13. Bausewein C, Daveson B, Benalia H, Simon ST, Higginson IJ. Outcome measurement in palliative care. London: PRISMA; 2011. Available from: <http://pos-pal.org/Resources.php>.
14. Götze H, Brähler E, Gansera L, Polze N, Köhler N. Psychological distress and quality of life of palliative cancer patients and their caring relatives during home care. *Support Care Cancer*. 2014;22(10):2775–82.
15. User's Guide for Implementing Patient-Reported Outcomes Assessment in Clinical Practice. Available from: <http://www.isoqol.org/UserFiles/file/UsersGuide.pdf> visited on the 27th July 2014.
16. Antunes B, Harding R, Higginson IJ, on behalf of EUROIMPACT. Implementing patient-reported outcome measures in palliative care clinical practice: a systematic review of facilitators and barriers. *Palliat Med*. 2014;28(2):158–75.
17. Etkind SN, Daveson BA, Kwok W, Witt J, Bausewein C, Higginson IJ, et al. Capture, transfer, and feedback of patient centred outcomes data in palliative care populations: does it make a difference? A systematic review. *J Pain Symptom Manag*. 2015;49(3):611–24. doi:10.1016/j.jpainsymman.2014.07.010
18. Murtagh F, Ramsenthaler C, Firth A, Groeneveld EI, Lovell N, Simon S et al. A brief, patient- and proxy-reported outcome measure for the adult palliative care population: validity and reliability of the Integrated Palliative Outcome Scale (IPOS) 9th world research congress of the European Association for Palliative Care (EAPC); 2016.
19. Antunes B, Ferreira PL. Integrated palliative care outcome scale: protocol validation for the portuguese population. *Rev Cuidados Paliat*. 2017;4:65–102.
20. Antunes B, Murtagh F, Bausewein C, Harding R, Higginson IJ. Screening for depression in advanced disease: psychometric properties, sensitivity and specificity of two items of the palliative care outcome scale (POS). *J Pain Symptom Manag*. 2015;49(2):277–88. doi:10.1016/j.jpainsymman.2014.06.014
21. Antunes B, Rodrigues PP, Higginson IJ, Ferreira PL. Validation and cultural adaptation of the integrated palliative care outcome scale (IPOS) for the Portuguese population. Poster at 5th world congress of the European Association for Palliative Care in Madrid, Spain on 18–20; 2017.
22. Antunes B, Rodrigues PP, Higginson IJ, Ferreira PL. Validation of the integrated palliative care outcome scale (IPOS) to the Portuguese population - completion assessment and thematic analysis of the open question items. Oral presentation at the 5th world congress of the European Association for Palliative Care in Madrid, Spain on 18–20 2017.
23. Pais-Ribeiro J, Silva I, Ferreira T, Martins A, Meneses R, Baltar M. Validation study of a Portuguese version of the hospital anxiety and depression scale. *Psychol Health Med*. 2007;12(2):225–37. doi:10.1080/13548500500524088
24. Acuña E, Rodriguez C. The treatment of missing values and its effect on classifier accuracy. In: Banks D, House L, FR MM, et al, eds. *Classification, Clustering and Data Mining Applications*. Berlin: Springer; 2004. 639e648 p.
25. Mistry B, Bainbridge D, Bryant D, Tan Toyofuku S, Seow H. What matters most for end-of-life care? Perspectives from community-based palliative care providers and administrators. *BMJ Open*. 2015;5(6):e007492. doi:10.1136/bmjopen-2014-007492
26. Wang T, Molassiotis A, Chung BPM, Tan JY. Unmet care needs of advanced cancer patients and their informal caregivers: a systematic review. *BMC Palliat Care*. 2018;17(1):96. doi:10.1186/s12904-018-0346-9
27. Rayner L, Price A, Hotopf M, Higginson IJ. The development of evidence-based European guidelines on the management of depression in palliative cancer care. *Eur J Cancer*. 2011;47(5):702–12.

28. Moens K, Higginson IJ, Harding R. EURO IMPACT. Are there differences in the prevalence of palliative care-related problems in people living with advanced cancer and eight non-cancer conditions? A systematic review. *J Pain Symptom Manag.* 2014;48(4):660–77.
29. Stiel S, Matthies DM, Seuss D, Walsh D, Lindena G, Ostgathe C. Symptoms and problem clusters in cancer and non-cancer patients in specialized palliative care—is there a difference? *J Pain Symptom Manag.* 2014;48(1):26–35.
30. Hotopf M, Chidgey J, Addington-Hall J, Ly KL. Depression in advanced disease: a systematic review. Part 1. Prevalence and case finding. *Palliat Med.* 2002;16(2):81-97.
31. Mitchell AJ. Are one or two simple questions sufficient to detect depression in cancer and palliative care? A Bayesian meta-analysis. *Br J Cancer.* 2008;98(12):1934–43. doi:10.1038/sj.bjc.6604396
32. Payne A, Barry S, Creedon B, Stone C, Sweeney C, O'Brien T, et al. Sensitivity and specificity of a two-question screening tool for depression in a specialist palliative care unit. *Palliat Med.* 2007;21(3):193-8.