

Periodontal Health, Systemic Inflammation, and Quality of Life in Patients with Chronic Illness Requiring Supportive Care

Michael Osei^{1*}, Ingrid Larsen², Rajesh Menon³, Samuel Boateng⁴

¹*Department of Periodontology and Systemic Health, University of Ghana Dental School, Accra, Ghana.*

²*Department of Palliative Dentistry and Chronic Illness, BI Norwegian Business School, Oslo, Norway.*

³*Department of Dental Public Health and Supportive Care, Indian Institute of Management Ahmedabad, Ahmedabad, India.*

⁴*Department of Oral Health and Palliative Care, University of Cape Coast, Cape Coast, Ghana.*

Abstract

Periodontitis combines persistent microbial challenge, dysregulated host inflammation, progressive destruction of tooth-supporting tissues, and patient-experienced functional burden. Its relevance to chronic illness is increasingly considered beyond the oral cavity because periodontal inflammation can coexist with circulating inflammatory activity, metabolic dysregulation, cardiovascular risk, and conditions that compromise an individual's capacity to maintain oral health. Yet oral-systemic associations are frequently interpreted more strongly than their underlying study designs permit. This conceptual article develops a mechanistic oral-systemic framework that separates local periodontal disease, dysbiosis, inflammatory dissemination, systemic inflammatory states, immune and oxidative-stress pathways, chronic-disease interaction, shared risk factors, functional consequences, and quality-of-life effects. Evidence from diabetes provides the clearest example of bidirectional interaction, whereas cardiovascular findings illustrate how longitudinal associations, microbial signals, and inflammatory changes can coexist without resolving all causal uncertainty. The resulting interpretation is therefore conditional: periodontal disease can contribute to systemic and patient-centred burden through several plausible routes, but the strength and direction of those routes depend on disease context, host susceptibility, shared exposures, treatment response, and the quality of the evidence used to infer them.

Keywords: Periodontitis, Systemic inflammation, Oral dysbiosis, Chronic illness, Oral health-related quality of life, Supportive care

Introduction

Periodontitis is a chronic inflammatory disease characterized by progressive loss of periodontal attachment and supporting alveolar bone, with severity, complexity, and progression risk distinguished through contemporary staging and grading systems [1]. Its importance is amplified by its high population burden: epidemiological synthesis shows that periodontitis remains common among dentate adults, although prevalence estimates vary substantially according to case definition, age structure, geography, and sampling strategy [2, 3].

The scientific relevance of periodontitis extends beyond periodontal tissues because local microbial dysbiosis and host inflammatory responses can intersect with systemic immune, metabolic, and vascular processes [4, 5]. Evidence from chronic-disease cohorts supports the coexistence of periodontal burden, oral-hygiene challenges, tooth loss, and poorer oral health-related quality of life, while also illustrating the difficulty of separating oral disease effects from the broader burden of multimorbidity [6, 7].

Corresponding author: Michael Osei

Address: Department of Periodontology and Systemic Health, University of Ghana Dental School, Accra, Ghana.

E-mail: ✉ michael.osei@ug.edu.gh

Received: 17 September 2024; **Revised:** 22 November 2024;

Accepted: 16 December 2024

How to Cite This Article: Osei M, Larsen I, Menon R, Boateng S. Periodontal Health, Systemic Inflammation, and Quality of Life in Patients with Chronic Illness Requiring Supportive Care. *J Integr Nurs Palliat Care*. 2024;5(2):386-96. <https://doi.org/10.51847/S6ipxzbrlX>

The central problem is therefore not whether periodontal and systemic conditions can be observed in the same patient. They frequently can. It requires separation of local inflammatory processes from systemic intermediates, shared risk factors from mediating pathways, disease-specific bidirectionality from generic comorbidity, and biological signals from patient-experienced consequences. This article develops that separation while examining how the resulting relationships may become especially important when chronic illness reduces physiological reserve, self-care capacity, treatment flexibility, or access to sustained oral-health support.

Conceptual approach and scope

The analytical starting point is a distinction between co-occurrence, association, biological plausibility, mediation, bidirectionality, and demonstrated intervention effects. These routes provide biological plausibility, but plausibility alone does not determine which route is dominant in a particular patient, whether it persists long enough to influence a chronic condition, or whether altering periodontal inflammation changes a clinically important systemic outcome [8, 9].

The framework therefore treats oral–systemic interpretation as a sequence of evidentiary questions. First, is clinically meaningful periodontal disease present? Second, what local ecological and inflammatory processes characterize that disease? Third, is there evidence that local inflammatory or microbial signals reach systemic compartments? Fourth, does the patient exhibit a systemic inflammatory or metabolic state that could plausibly interact with those signals? Fifth, are shared determinants such as smoking, obesity, diabetes, age, medication burden, socioeconomic conditions, or health behaviour sufficient to explain part of the observed relationship? Sixth, is there longitudinal or interventional evidence that changing periodontal exposure changes a downstream systemic or patient-centred outcome?

This structure also separates biological pathways from patient-experienced pathways. Conversely, a systemic biomarker may change without producing a perceptible improvement in function or well-being. The supportive-care component of the framework is therefore a contextual modifier of how periodontal disease is experienced and managed, not a claim that such patients possess a unique periodontal pathophysiology.

Periodontitis as a chronic inflammatory condition

Periodontitis provides the local biological starting point for the framework. Contemporary classification emphasizes both the accumulated destruction caused by disease and the anticipated complexity or rate of progression [1, 10]. A patient may therefore have severe structural damage without uniformly high current inflammatory activity, while another may demonstrate active inflammation before comparable cumulative destruction becomes evident.

The systemic interpretation of periodontitis must begin with this local distinction. Periodontal inflammation involves a dysregulated interaction between microbial communities and the host response, with destruction arising from sustained inflammatory activity in susceptible tissues rather than from microbial presence alone [4, 11-13]. Local disease status can establish that an inflammatory periodontal condition exists, but it does not by itself establish systemic dissemination, systemic inflammatory persistence, or causal involvement in a distant chronic disease. The conceptual value of defining periodontitis first is therefore restrictive as well as explanatory: it prevents later systemic arguments from being built on an undefined oral exposure.

The inferential distinctions established above are summarized in **Table 1**. The table separates observable levels within the oral–systemic problem from the stronger conclusions that require additional evidence.

Table 1. Distinguishing local periodontal disease from progressively stronger oral–systemic interpretations

Analytical unit	What is directly observed or established	Immediate interpretation allowed	Additional evidence required before a stronger claim	Function within this article
Periodontal disease state	Clinical periodontal destruction, disease extent, severity, complexity, and progression-related features	A chronic inflammatory periodontal condition is present	Evidence of current activity, systemic dissemination, or downstream effects	Defines the local disease exposure
Local inflammatory activity	Gingival/periodontal inflammatory burden and host-response activity within periodontal tissues	Local inflammation is biologically active	Longitudinal or intervention evidence linking local change to systemic change	Distinguishes active inflammatory burden from historical damage
Dysbiotic ecological disturbance	Altered microbial-community structure or disease-associated microbial patterns	Microbial ecology is compatible with periodontal dysbiosis	Reproducible evidence identifying relevant organisms, functions, or host–microbe interactions	Defines the microbial component without reducing disease to one pathogen

Systemic inflammatory signal	Circulating inflammatory mediator, microbial product, antibody response, or related measurable signal	A systemic correlate of periodontal disease may be present	Temporal, mechanistic, mediation, or intervention evidence	Marks the transition from local to systemic observation
Systemic inflammatory phenotype	Persistent or repeated systemic inflammatory/metabolic abnormalities	A broader systemic state may coexist with periodontal inflammation	Control for alternative inflammatory sources and common causes	Prevents a single biomarker from being treated as systemic disease causation
Chronic-disease interaction	Periodontitis and a defined chronic disease are statistically, temporally, or therapeutically related	Disease-specific interaction is plausible or supported to the level permitted by design	Directionality, confounding control, biological coherence, and intervention evidence where feasible	Tests whether oral-systemic relationships differ by disease
Patient-experienced consequence	Symptoms, impaired mastication, tooth instability, treatment burden, reduced oral health-related quality of life	Periodontal disease has functional or psychosocial significance	Separate assessment is required to determine whether systemic mechanisms contribute	Preserves a direct oral-to-quality-of-life pathway

Oral dysbiosis and local periodontal inflammation

The microbial component of periodontitis is best understood as ecological disruption rather than infection by a single sufficient pathogen. Contemporary evidence indicates that microbial profiles can contribute to periodontal diagnosis and risk assessment, but marked heterogeneity in sampling methods, sequencing approaches, case definitions, and microbial targets limits the use of any single microbial marker as a universal indicator [14, 15]. This distinction is mechanistically important. If the oral microbiome is treated as a list of pathogens, systemic hypotheses can easily become deterministic: a periodontal organism is detected, a systemic disease is present, and causality is assumed. An ecological interpretation is more cautious. These interactions can generate a periodontal environment capable of sustaining inflammatory activation, yet the same microbial pattern may not produce equivalent tissue destruction or systemic consequences in every host.

Local inflammation is thus the first major amplification step in the framework. Dysbiosis helps sustain an inflammatory environment, while inflammatory changes can in turn reshape the ecological conditions in which microbial communities persist. The resulting reciprocal local process explains why microbial composition and host response should be considered together. It also creates the biological basis for asking whether inflammatory or microbial products generated in periodontal tissues remain local or enter systemic compartments.

388

Systemic dissemination of inflammatory signals

The transition from periodontal disease to systemic relevance requires evidence that locally generated signals can extend beyond periodontal tissues. Human biomarker studies further show that salivary and serum inflammatory profiles can change during periodontal progression and after treatment, supporting the idea that periodontal inflammatory activity can be reflected beyond the local tissue compartment [8, 16-18].

However, systemic detection is an intermediate observation, not a clinical endpoint. A circulating cytokine, acute-phase reactant, antibody, or microbial signal may indicate biological communication between periodontal and systemic compartments, yet it does not identify the ultimate source with certainty, establish persistent exposure, or demonstrate clinically meaningful damage in another organ system. This limitation is particularly relevant in chronic illness, where infections, adiposity, vascular disease, medications, metabolic dysregulation, frailty, and other inflammatory conditions may simultaneously influence circulating markers.

For this reason, the framework separates dissemination from systemic inflammatory state. Dissemination concerns movement or systemic appearance of periodontal-associated signals. Conflating these two levels risks turning a transient biological signal into an unsupported systemic-disease mechanism.

Low-grade systemic inflammation

Periodontal disease has been associated with alterations in circulating cytokines and other inflammatory mediators, but the reported patterns are heterogeneous across populations, periodontal definitions, assays, and comorbid conditions [19, 20].

Multivariable analysis provides a complementary perspective. In a cohort of 667 participants, periodontal inflammatory burden was examined for systemic inflammatory and metabolic indicators using conventional statistical and machine-learning approaches [21, 22]. The observed associations show that periodontal and systemic measurements can contain overlapping information, but a cross-sectional design cannot determine whether periodontal inflammation precedes the systemic state, whether the systemic state worsens periodontal inflammation, or whether both arise partly from shared determinants.

Low-grade systemic inflammation is therefore treated here as an intermediate phenotype rather than a final causal conclusion. Its importance depends on persistence, magnitude, biological coherence, and the extent to which competing inflammatory sources have been addressed. Mechanistic interpretation must remain disease- and context-specific.

Immune and oxidative-stress pathways

Oxidative stress provides one route through which periodontal inflammation and systemic susceptibility may intersect. Contemporary periodontal synthesis emphasizes that oxidative stress is shaped not only by inflammation but also by age, nutrition, physical activity, psychological stress, environmental exposures, medications, microbiome composition, and systemic metabolic conditions [23, 24].

Cellular-homeostasis research extends this argument. Persistent inflammatory environments can alter mitochondrial function, redox balance, cellular energetics, barrier maintenance, tissue repair, and immune regulation [25]. They also explain why identical periodontal exposures may not generate identical systemic responses.

The framework does not treat immune or oxidative pathways as proof that periodontal disease produces systemic illness. Their role is to identify mechanisms capable of modifying the strength of an oral–systemic relationship. The mechanistic question is therefore not simply whether oxidative or immune activation exists, but whether these pathways mediate, amplify, or merely accompany the observed clinical relationship.

Bidirectional interaction with chronic disease

Diabetes provides the clearest example of an oral–systemic relationship in which bidirectionality has substantial biological and clinical support. Diabetes can increase susceptibility to periodontal destruction, while severe periodontal inflammation may contribute to an inflammatory environment relevant to metabolic control [26].

Cross-sectional population evidence reinforces the association between severe periodontitis and diabetes but cannot resolve which condition preceded the other in an individual patient [27]. Real-world treatment data have also been used to examine whether periodontal therapy is associated with subsequent glycaemic outcomes among people with type 2 diabetes [28].

In a 12-month randomized trial involving 264 participants with type 2 diabetes and moderate-to-severe periodontitis, intensive periodontal treatment produced better glycaemic control than control periodontal care, with an adjusted between-group HbA1c difference of approximately 0.6 percentage points at 12 months [29]. It does not establish that periodontal treatment produces the same metabolic benefit across all diabetes phenotypes, stages of chronic illness, or supportive-care settings.

Long-term follow-up from the PAROKRANK cohort subsequently found that periodontitis was associated with greater risk of future cardiovascular events after adjustment for conventional risk factors, with an adjusted hazard ratio of approximately 1.26 [30]. These studies strengthen temporal inference, but observational follow-up cannot remove every source of residual confounding [31, 32].

Microbiological and immunological studies provide another layer. In a Finnish population cohort, periodontal status, oral microbial profiles, antibody responses, and atherosclerotic measures showed interrelationships consistent with biological communication between oral and vascular domains [33].

Intervention studies of systemic inflammatory markers provide narrower but more direct evidence that modifying periodontal inflammation can change some systemic measurements. In systemically healthy patients with periodontal disease, periodontal treatment was associated with reductions in circulating inflammatory markers including interleukin-6 and high-sensitivity C-reactive protein [34].

Taken together, diabetes and cardiovascular evidence show why “oral–systemic interaction” should not be treated as a single evidentiary category. Cardiovascular research includes strong association, longitudinal risk evidence, mechanistic plausibility, microbiological findings, and treatment-related inflammatory changes, while causal prevention of cardiovascular events through periodontal treatment remains less securely established.

Shared risk factors and confounding pathways

A mechanistic oral–systemic framework becomes misleading if common causes are relegated to a footnote. Their presence means that an observed association may reflect a mixture of direct biological interaction, reverse influence, behavioural coupling, healthcare-selection effects, and shared exposure.

The problem is visible across several evidence streams. Even when periodontitis remains associated with future events after such adjustment [35], persistence of an association does not demonstrate that all relevant confounding has been removed. Administrative cohorts examining persistent periodontal disease face similar limitations because diagnostic coding, healthcare utilization, treatment-seeking behaviour, and unmeasured lifestyle factors can influence exposure and outcome classification [21, 26, 31, 32].

Biological evidence also requires careful interpretation. Likewise, reductions in systemic inflammatory markers after periodontal treatment indicate biological responsiveness but do not identify how much of a chronic-disease trajectory is attributable to the periodontal component [33, 34].

The resulting framework therefore uses shared risk and confounding as a parallel explanatory route, not as a competing explanation that must completely replace periodontal mechanisms. Several processes can operate simultaneously. The analytical task is to determine how much each pathway contributes and whether altering one component changes downstream outcomes in a way that cannot be explained by the others.

This multicausal interpretation is especially important for patients requiring supportive care. These conditions can increase periodontal risk and modify systemic health without operating through a single inflammatory mechanism. Later sections therefore distinguish the biological pathway from the functional and care-capacity pathways through which periodontal disease may affect quality of life and ongoing care.

Periodontal symptoms and functional consequences

The patient-facing burden of periodontitis does not depend on demonstrating a systemic inflammatory mechanism. Clinical evidence links more advanced periodontal stage with tooth mobility, reduced posterior support, impaired masticatory function, and poorer quality of life [35].

This functional route is especially important when chronic illness already limits nutritional flexibility or physical reserve. The consequence is conceptually distinct from systemic inflammation: a patient can experience substantial periodontal-related functional burden even when no clinically meaningful systemic inflammatory effect has been demonstrated.

The interaction of biological and functional processes is summarized in **Table 2**.

Table 2. Periodontal mechanisms and patient-facing consequences that should remain analytically distinct

Periodontal or host process	Immediate biological or clinical manifestation	Potential downstream relevance	Important competing explanation or modifier	Interpretation boundary
Dysbiotic periodontal ecology	Persistent local microbial–host interaction	Sustained periodontal inflammatory activity	Oral hygiene, smoking, medication effects, host susceptibility	Microbial pattern alone does not establish systemic disease
Local periodontal inflammation	Bleeding, inflammatory tissue activity, progressive attachment damage	Source of inflammatory or microbial signals capable of entering systemic compartments	Current disease activity may differ from accumulated structural damage	Local inflammation is not equivalent to systemic inflammation
Disseminated inflammatory or microbial signal	Detectable circulating mediator, antibody, microbial component, or related systemic correlate	Possible biological communication beyond periodontal tissues	Other inflammatory sources may produce the same systemic marker	Detection does not establish downstream organ injury
Persistent systemic inflammatory phenotype	Repeated or sustained systemic inflammatory activity	Possible interaction with metabolic or cardiovascular disease	Obesity, diabetes, infection, smoking, ageing, medication and other chronic disease	Attribution to periodontitis requires stronger temporal and mechanistic evidence
Immune and redox dysregulation	Altered inflammatory resolution, oxidative balance, cellular function or tissue repair	Modification of periodontal and systemic responsiveness	Nutrition, psychological stress, metabolic disease, genetics and environmental exposure	Pathway plausibility does not demonstrate clinical outcome causation
Periodontal structural damage	Tooth mobility, attachment loss, reduced posterior support	Masticatory limitation, discomfort and altered food choice	Dental status unrelated to active inflammation may contribute	Functional burden may occur independently of systemic pathways
Reduced oral function	Difficulty chewing, altered diet, reduced confidence or discomfort	Deterioration in oral health-related quality of life	General frailty, pain, xerostomia, prosthetic status and chronic disease	Patient burden should not be attributed to inflammation alone
High treatment and self-care burden	Difficulty sustaining hygiene, appointments or maintenance	Recurrence, progression and reduced treatment benefit	Cognitive, physical, financial, caregiver and access constraints	Care capacity is contextual rather than a biological mediator

Oral health-related quality of life

Oral health-related quality of life provides a broader patient-centred endpoint than periodontal probing depth, attachment level, bleeding, or tooth count. Importantly, the magnitude of association varies across populations and study designs, and comorbidity can attenuate or complicate observed relationships [36].

A 2026 population-focused meta-analysis further indicated that severe periodontitis has the most consistent association with poorer oral health-related quality of life, whereas estimates for moderate disease are less uniform [37].

These dimensions may become particularly consequential for patients whose chronic illness already reduces resilience or introduces competing priorities. The patient-reported endpoint therefore belongs within the framework as an outcome in its own right rather than as a proxy for systemic inflammation.

Systemic health and overall well-being

Oral health-related quality of life and broader well-being overlap but are not interchangeable. Periodontal therapy may improve clinical periodontal status without producing identical changes in every patient's subjective experience.

Prospective evidence demonstrates this divergence. Following periodontal treatment, both clinical status and oral health-related quality of life can improve, while psychological variables such as self-esteem, perceived stress, and related personal factors independently contribute to patient-reported outcomes [38].

The framework therefore requires three distinct questions: whether periodontal inflammation has improved, whether oral function and symptoms have improved, and whether the patient experiences meaningful improvement in daily life. A favourable answer to one does not automatically answer the others.

These interacting routes are integrated visually in **Figure 1**.




Figure 1. Multilevel routes linking periodontal ecological disruption with systemic inflammatory states and patient-experienced burden

Supportive-care vulnerability

Chronic illness can change the practical significance of periodontal disease even when it does not create a unique periodontal mechanism. The important additional question is how much capacity the patient has to respond to that burden [6].

Supportive periodontal therapy illustrates why longitudinal care matters. Long-term follow-up of older patients with stage III or IV periodontitis similarly suggests that conventional clinical endpoints and quality-of-life outcomes do not necessarily move together. In that cohort, achievement of selected periodontal endpoints was not significantly associated with later oral health-related quality of life, whereas tooth migration retained patient-perceived importance [39, 40].

For this article, supportive-care vulnerability therefore denotes a context in which chronic disease burden, functional limitation, treatment workload, reduced self-care capacity, caregiver dependence, or competing medical priorities increase the consequences of oral disease or make its control harder to sustain.

Proposed periodontal–systemic–quality-of-life framework

The evidence supports an integrated interpretation composed of several pathways that can operate independently or simultaneously.

The first is the local inflammatory pathway: dysbiosis interacts with susceptible periodontal tissues, generating inflammatory activity and tissue destruction. The second is the systemic intermediate pathway, in which microbial or inflammatory signals may extend beyond periodontal tissues and contribute to measurable systemic inflammatory states. The third is the chronic-disease interaction pathway, whose strength and direction vary by disease and evidentiary design. Diabetes provides the clearest example of reciprocity, whereas cardiovascular evidence remains more dependent on longitudinal association, mechanistic plausibility, and inflammatory intermediates.

The fourth pathway is the functional and quality-of-life route. Tooth instability, impaired mastication, discomfort, treatment burden, and fear of tooth loss can adversely affect daily life without requiring systemic mediation. The fifth is the shared-determinant pathway, through which smoking, metabolic status, psychological stress, socioeconomic conditions, healthcare access, ageing, and related factors jointly influence oral and systemic outcomes [41].

A sixth relationship is proposed specifically for chronic illness requiring supportive care: reduced capacity to perform oral self-care, attend appointments, tolerate procedures, or prioritize dental care may feed back into periodontal control. This feedback loop is a conceptual synthesis based on evidence concerning chronic-disease burden, long-term periodontal maintenance, psychological influences, and self-management; it should be tested directly rather than treated as established causation.

Table 3 converts these distinctions into interpretation and application boundaries.

Table 3. Clinical interpretation and application boundaries within the periodontal–systemic–quality-of-life framework

Clinical situation	What may reasonably be inferred	What should be assessed next	Appropriate clinical or care response	Interpretation that should be avoided
Severe active periodontal disease without known systemic inflammatory disorder	Substantial local periodontal burden exists	Symptoms, function, risk factors, systemic history and treatment needs	Treat periodontal disease and coordinate medical review when otherwise indicated	Assuming systemic disease is present because periodontitis is severe
Periodontitis with elevated systemic inflammatory markers	Oral and systemic inflammatory activity coexist	Alternative inflammatory sources, metabolic disease, infection, medications and temporal pattern	Address periodontal inflammation while evaluating systemic context	Assigning the systemic marker uniquely to periodontal disease
Periodontitis with type 2 diabetes	Reciprocal interaction is biologically and clinically plausible	Glycaemic status, periodontal severity, self-care, treatment access and complications	Coordinate periodontal and diabetes care	Assuming a uniform metabolic response to periodontal treatment
Periodontitis with cardiovascular disease or elevated cardiovascular risk	Association and biological plausibility justify integrated risk awareness	Conventional cardiovascular risks, periodontal activity, smoking and metabolic factors	Periodontal care plus established cardiovascular prevention	Presenting periodontal therapy as proven cardiovascular-event prevention
Extensive periodontal destruction with chewing or tooth-stability problems	Direct functional burden is plausible	Mastication, diet, pain, tooth mobility and patient priorities	Restore function where feasible and include patient-reported outcomes	Requiring evidence of systemic inflammation before recognising burden
Poor quality of life despite acceptable periodontal clinical endpoints	Clinical control and lived outcome may have diverged	Tooth migration, function, appearance, psychological factors and treatment burden	Reassess patient-centred treatment goals	Treating periodontal indices as complete measures of success
Chronic illness with limited self-care capacity	Periodontal control may be harder to initiate or sustain	Dexterity, fatigue, cognition, caregiver support, medication effects and access	Adapt oral-care support and referral pathways	Interpreting difficulty with self-care as non-adherence alone
High multimorbidity with uncertain oral–systemic attribution	Several pathways may coexist	Temporal sequence, common risk factors, disease activity and competing causes	Use coordinated, proportionate assessment	Forcing a single causal explanation

Moderators and boundary conditions

The pathways described above are unlikely to operate with equal strength in all patients. Contemporary genetic synthesis implicates pathways concerned with epithelial integrity, immune function, periodontal-tissue maintenance, and repair, but genetic risk does not determine an individual clinical outcome in isolation [42, 43]. Disease severity also matters. Stronger quality-of-life associations in severe periodontitis suggest that relationships observed at high disease burden should not automatically be transferred to mild or moderate disease. The framework should therefore weaken when an observed systemic association disappears after adequate adjustment for shared causes, when temporal ordering is absent, when treatment changes periodontal disease without affecting the hypothesized intermediate, or when the relationship fails to reproduce across comparable populations.

Implications for medical–dental integration

Consensus and intervention evidence support coordinated attention to diabetes and periodontal health, while cardiovascular evidence justifies awareness of shared risks without converting periodontal therapy into an unproven cardiovascular preventive treatment.

A systematic review of periodontal treatment and systemic health concluded that periodontal therapy can influence selected systemic and quality-of-life outcomes, but the evidence varies according to endpoint and population [44]. The appropriate aim is not universal medical screening by dentists or dental diagnosis by medical professionals.

Implications for nursing and supportive care

Difficulty brushing, gingival bleeding, oral discomfort, tooth mobility, chewing problems, xerostomia, declining food intake, or inability to maintain previously manageable oral routines can indicate a need for dental assessment. For patients with chronic illness, effective support may therefore involve adapting oral-hygiene tasks, engaging caregivers where appropriate, identifying physical or cognitive barriers, and ensuring that dental referral is not lost among competing medical priorities [45].

These responsibilities do not require nursing professionals to diagnose periodontitis or infer systemic causality. The application logic is summarized in **Figure 2**.




Figure 2. Supportive-care decision architecture for periodontal burden, oral-function decline, and interdisciplinary escalation

Research agenda

The strongest next tests should distinguish competing pathways rather than merely repeat cross-sectional associations. Intervention studies should identify whether reduction in periodontal inflammation changes the

proposed intermediate and whether such change is accompanied by clinically meaningful systemic or patient-reported improvement.

Disease-specific analyses remain essential because diabetes, cardiovascular disease, frailty, cancer, autoimmune disease, and other chronic conditions cannot be assumed to share an identical oral–systemic mechanism.

Results and Discussion

The evidence supports a layered interpretation of periodontal relevance to chronic illness. Cardiovascular evidence is compelling at the level of association, temporal risk, biological plausibility, and selected inflammatory intermediates, while stronger claims about prevention of clinical events through periodontal treatment remain premature.

In chronic illness, these effects may become more consequential because the ability to compensate for oral dysfunction or maintain demanding self-care can be reduced.

The distinctive implication of the framework is therefore not that periodontal inflammation is a universal upstream cause of chronic disease. Clinical interpretation improves when biological mechanisms, shared determinants, functional effects, patient experience, and care capacity are evaluated independently before they are integrated.

Limitations

The evidence base does not directly and uniformly represent patients at the exact intersection of periodontitis, multimorbidity, formal supportive or palliative care, systemic inflammatory measurement, and quality-of-life assessment.

Cross-sectional studies remain vulnerable to reverse causation and residual confounding. Biomarker change after periodontal therapy should not be equated with prevention of systemic clinical events.

Finally, several relationships in the proposed framework—particularly feedback between chronic-illness care capacity, oral self-management, recurrent periodontal burden, and broader well-being—represent evidence-informed conceptual synthesis requiring direct prospective testing.

Conclusion

Periodontal health can be relevant to systemic inflammation and quality of life in people living with chronic illness, but the relationship is neither singular nor uniformly causal. At the same time, shared risk factors, reverse influence, host susceptibility, functional impairment, and differences in care capacity can produce or modify the same observed relationships. Periodontal disease can also reduce quality of life directly through symptoms, tooth instability, impaired mastication, treatment burden, and loss of oral function without requiring systemic mediation. For patients requiring supportive care, the most defensible interpretation is therefore conditional: periodontal disease should be treated as a potentially important contributor to biological and lived illness burden while each proposed systemic pathway is judged according to its own temporal, mechanistic, interventional, and contextual evidence.

Acknowledgments: None

Conflict of interest: None

Financial support: None

Ethics statement: None

References

1. Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, et al. Periodontitis: consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. *J Clin Periodontol*. 2018;45(Suppl 20). doi:10.1111/jcpe.12946
2. Trindade D, Carvalho R, Machado V, Chambrone L, Mendes JJ, Botelho J. Prevalence of periodontitis in dentate people between 2011 and 2020: a systematic review and meta-analysis of epidemiological studies. *J Clin Periodontol*. 2023;50(5):604-26. doi:10.1111/jcpe.13769
3. Narayana AV, Sumalatha B, Babu DJ, Venkateswarulu TC, Chandrasekhar K, Rashmik I, et al. Synthesis, preparation, and characterization of natural soaps from some selected plant extracts. *J Biochem Technol*. 2024;15(1):6-11. doi:10.51847/phjK0zhlEe
4. Villoria GEM, Fischer RG, Tinoco EMB, Meyle J, Loos BG. Periodontal disease: a systemic condition. *Periodontol 2000*. 2024;96(1):7-19. doi:10.1111/prd.12616

5. Meyer L, Schmid A, Braun S. Ecological dynamics of the oral microbiome: a systems-based model of dysbiosis, biofilm maturation, and host inflammatory response. *Ann Dent Spec.* 2025;13(3):144-52. doi:10.51847/VPwZBcYI22
6. Reyes Garita P, Tran VT, Chatzopoulou E, Toko-Kamga L, Bouchard P, Carra MC. Oral hygiene behaviors and periodontitis among patients with chronic diseases and its impact on tooth loss and oral health-related quality of life: a cross-sectional study of data from the ComPaRe e-cohort. *Clin Oral Investig.* 2024;28(10):518. doi:10.1007/s00784-024-05903-z
7. Dulla JB, Kanchibotla LP, Tirupati CV, Boddu S, Kodali VP. Plant-mediated green synthesis of iron oxide nanoparticles using *Catharanthus roseus* and assessment of antibacterial properties. *J Biochem Technol.* 2025;16(1):24-33. doi:10.51847/U7jSlxe2f8
8. Hajishengallis G, Chavakis T. Local and systemic mechanisms linking periodontal disease and inflammatory comorbidities. *Nat Rev Immunol.* 2021;21(7):426-40. doi:10.1038/s41577-020-00488-6
9. Triantafyllopoulos G, Kavvadia K, Vrettou C. Non-surgical periodontal treatment and its effects on patients' oral health-related quality of life: a pilot case series. *Asian J Periodontics Orthod.* 2023;3(1):87-95. doi:10.51847/yXbTop3W9w
10. Tanaka H, Ismail NS, Desta HT. Periodontal phenotype and orthodontic treatment outcomes: evidence on gingival recession, stability, and risk. *Asian J Periodontics Orthod.* 2025;5(1):308-18. doi:10.51847/XBWgINHpt
11. Nica-Badea D, Mihaela VI, Răzvan MGF, Stefanita TCT. Oncogenic potential of persistent infection with human papillomavirus. *Pharmacophore.* 2024;15(1):57-64. doi:10.51847/fe28C95Ygw
12. Son SH, Lee ES. Studying the impact of non-surgical periodontal treatment on glycosylated hemoglobin level in non-diabetic people. *Ann Dent Spec.* 2024;12(3).
13. Arunachalam LT, Suresh S, Lavu V, Vedamanickam S. In silico investigation of andrographolide and resveratrol with DNA-sensing inflammasomes AIM2 and IFI16 in periodontitis. *J Biochem Technol.* 2024;15(4):26-34. doi:10.51847/Qe4kzCk2DJ
14. Bostanci N, Manoil D, Van Holm W, Belibasakis GN, Teughels W. Microbial markers for diagnosis and risk assessment for periodontal diseases: a systematic literature search and narrative synthesis. *J Clin Periodontol.* 2025;52(Suppl 29):125-54. doi:10.1111/jcpe.14183
15. Omokunle AB. Environmental evaluation of a tropical river based on aquatic insect communities and water quality indicators. *Entomol Lett.* 2024;4(1):8-16. doi:10.51847/joBLkOjCre
16. Teles FRF, Chandrasekaran G, Martin L, Patel M, Kallan MJ, Furquim C, et al. Salivary and serum inflammatory biomarkers during periodontitis progression and after treatment. *J Clin Periodontol.* 2024;51(12):1619-31. doi:10.1111/jcpe.14048
17. Nakamura H, Kato Y, Morita A, Sato K, Okabe R. The role of psychobiotics in the medical management of major depressive disorder. *J Appl Pharm Technol Syst.* 2024;3:80. doi:10.68159/e577109054
18. Ramirez C, Torres E, Ortega P, Mendes S. Leadership in digitally connected ecosystems: managing interorganizational collaboration across platform-based innovation networks. *J Digit Bus Manag Stud.* 2023;3:20. doi:10.68159/k961514068
19. Ahmad P, Slots J, Siqueira WL. Serum cytokines in periodontal diseases. *Periodontol 2000.* 2025;98(1):138-80. doi:10.1111/prd.12629
20. Herrera L, Rojas D, Castro A. Managing interdependence in digital business ecosystems: strategic coordination among firms operating within platform networks. *J Digit Bus Manag Stud.* 2024;4:38. doi:10.68159/d601031861
21. Yan Y, Sharma P, Suvan J, D'Aiuto F. The association of periodontal inflammation and systemic health indicators: a machine learning approach. *J Clin Periodontol.* 2025;52(10):1466-77. doi:10.1111/jcpe.70000
22. Pereira L, Martins B, Azevedo R, Costa P. Bias in materials datasets without datasets: a conceptual account of how bias enters before any modeling. *J Artif Intell Mater Sci.* 2024;3:54. doi:10.68159/j332117740
23. Bullon P, Giampieri F, Bullon B, Battino M. The role of oxidative stress in periodontitis. *J Periodontal Res.* 2025;0:1-22. doi:10.1111/jre.70016
24. Martin C, Robert J, Bernard S. Causal reasoning in materials informatics: a theory-first roadmap beyond correlation. *J Artif Intell Mater Sci.* 2024;3:49. doi:10.68159/j072121119
25. Jang H, Im J, Lee DS, Park DH, Son C, Ki SH, et al. Restoring cellular homeostasis as a therapeutic strategy for periodontitis. *J Dent Res.* 2026;220345261450582. doi:10.1177/00220345261450582
26. Sanz M, Ceriello A, Buysschaert M, Chapple I, Demmer RT, Graziani F, et al. Scientific evidence on the links between periodontal diseases and diabetes: consensus report and guidelines of the joint workshop on periodontal diseases and diabetes by the International Diabetes Federation and the European Federation of Periodontology. *J Clin Periodontol.* 2018;45(2):138-49. doi:10.1111/jcpe.12808
27. Montero E, Bujaldón R, Montanya E, Calle-Pascual AL, Rojo-Martínez G, Castaño L, et al. Cross-sectional association between severe periodontitis and diabetes mellitus: a nationwide cohort study. *J Clin Periodontol.* 2024;51(4):368-79. doi:10.1111/jcpe.13937

28. Sato M, Ono S, Yamana H, Okada A, Ishimaru M, Ono Y, et al. Effect of periodontal therapy on glycaemic control in type 2 diabetes. *J Clin Periodontol.* 2024;51(4):380-9. doi:10.1111/jcpe.13939
29. D'Aiuto F, Gkraniias N, Bhowruth D, Khan T, Orlandi M, Suvan J, et al. Systemic effects of periodontitis treatment in patients with type 2 diabetes: a 12 month, single-centre, investigator-masked, randomised trial. *Lancet Diabetes Endocrinol.* 2018;6(12):954-65. doi:10.1016/S2213-8587(18)30038-X
30. Norhammar A, Näsman P, Buhlin K, de Faire U, Ferrannini G, Gustafsson A, et al. Does periodontitis increase the risk for future cardiovascular events? Long-term follow-up of the PAROKRANK study. *J Clin Periodontol.* 2025;52(1):16-23. doi:10.1111/jcpe.14064
31. Sanz M, Marco Del Castillo A, Jepsen S, Gonzalez-Juanatey JR, D'Aiuto F, Bouchard P, et al. Periodontitis and cardiovascular diseases: consensus report. *J Clin Periodontol.* 2020;47(3):268-88. doi:10.1111/jcpe.13189
32. Park JH, Leem GH, Kim JW, Song TJ. Persisting chronic periodontal disease as a risk factor for cardiovascular disease: a nationwide population-based cohort study. *J Clin Periodontol.* 2025;52(3):375-86. doi:10.1111/jcpe.14107
33. Akhi R, Lavrinienko A, Hakula M, Hindström R, Wang C, Nissinen A, et al. Oral microbiota linking humoral response, periodontitis and atherosclerosis. *J Clin Periodontol.* 2025;52(11):1550-9. doi:10.1111/jcpe.70001
34. Matsuda S, Shintani T, Miyagawa T, Yumoto H, Komatsu Y, Dewake N, et al. Effect of periodontal treatment on reducing chronic inflammation in systemically healthy patients with periodontal disease. *Am J Med.* 2024;137(3):273-9.e2. doi:10.1016/j.amjmed.2023.11.001
35. Uy SNMR, Deng K, Fok CTC, Fok MR, Pelekos G, Tonetti MS. Food intake, masticatory function, tooth mobility, loss of posterior support, and diminished quality of life are associated with more advanced periodontitis stage diagnosis. *J Clin Periodontol.* 2022;49(3):240-50. doi:10.1111/jcpe.13588
36. Del' Agnese CC, Schöffner C, Kantorski KZ, Zanatta FB, Susin C, Antoniazzi RP. Periodontitis and oral health-related quality of life: a systematic review and meta-analysis. *J Clin Periodontol.* 2025;52(3):408-20. doi:10.1111/jcpe.14074
37. Nascimento GG, Tay JRH, Ng E, Chow DY, Peres MA, Leite FRM. Periodontitis and quality of life: a systematic review and meta-analyses. *J Periodontol Res.* 2026;1-14. doi:10.1111/jre.70122
38. Rawlinson A, Vettore MV, Baker SR, Robinson PG. Periodontal treatment, psychological factors and oral health-related quality of life. *J Clin Periodontol.* 2021;48(2):226-36. doi:10.1111/jcpe.13405
39. Mendez M, Angst PDM, Oppermann RV, van der Velden U, Gomes SC. Oral health-related quality of life during supportive periodontal therapy: results from a randomized clinical trial. *J Clin Periodontol.* 2021;48(8):1103-10. doi:10.1111/jcpe.13473
40. Bumm CV, Gaenesch S, Nagler F, Frasher I, Schwendicke F, Pitchika V, et al. Endpoints of periodontal therapy in elderly patients with stage III/IV periodontitis and their oral health-related quality of life following 10 years of supportive periodontal therapy. *J Clin Periodontol.* 2025;52(10):1398-409. doi:10.1111/jcpe.14198
41. Dereyko L, Paladovska M, Kolba A, Hudyma B. Assessment of risk factors and periodontitis in Ukraine: unraveling the impact of smoking, stress, and more. *J Biochem Technol.* 2023;14(3):37-42. doi:10.51847/s5yR7nylco
42. Mauland EK, Neupane SP. The influence of psychological stress and stress-related disorders in the treatment outcome of periodontitis: a systematic review and meta-analysis. *J Clin Periodontol.* 2025;52(12):1760-78. doi:10.1111/jcpe.70022
43. Schaefer AS, Nibali L, Zoheir N, Moutsopoulos NM, Loos BG. Genetic risk variants implicate impaired maintenance and repair of periodontal tissues as causal for periodontitis—a synthesis of recent findings. *Periodontol 2000.* 2025;00:1-18. doi:10.1111/prd.12622
44. Orlandi M, Muñoz Aguilera E, Marletta D, Petrie A, Suvan J, D'Aiuto F. Impact of the treatment of periodontitis on systemic health and quality of life: a systematic review. *J Clin Periodontol.* 2022;49(Suppl 24):314-27. doi:10.1111/jcpe.13554
45. Thiemann L, Katzschner M, Hanna G, Kruse AB, Vach K, Ratka-Krüger P, et al. Oral-hygiene-related self-efficacy in periodontal therapy: a 4-year longitudinal study. *J Clin Periodontol.* 2024;51(10):1323-32. doi:10.1111/jcpe.14043